



Webmail
Univ. Bung Hatta

azrita ubh <azrita31@bunghatta.ac.id>

Manuscript 124706 conditionally accepted for publication

F1000.Research <research@f1000.com>
To: azrita ubh <azrita31@bunghatta.ac.id>

Wed, Nov 16, 2022 at 11:07 AM

Hi Azrita

I have been through the revised m/s – thanks.

There are a handful of text points that I need to raise with you, about the new changes, before I can put the m/s into production. I am putting these below, after the end of this email, with a reference to the place in the manuscript that the points apply to. You will see that I have suggested solutions or am double-checking a possible solution, in each case.

Also, there is a point about the Figures and their resolution that I need to raise, as this is unfortunately somewhat unresolved. Hopefully this will be easy to fix...

I look forward to hearing back on these final points, so we can get this interesting manuscript to the typesetters for printing.

Best regards

Jonathan

F1000 Research editorial team

Final points:

- re: *Figures query*. You say that Figures 1 and 2 are formatted at 300dpi. Perhaps the wrong versions have therefore been uploaded by mistake? 9or you have been passed the wrong version.) Please check this. According to our system (where the DPI shows up if you right-click on the picture file, scroll to Properties, and go to Details), the figures are showing as follows: Fig 1 - 150dpi; Fig 2 - 220dpi; Fig 3 - 300dpi; Fig 4 - 220dpi.

I'm afraid I cannot do anything about this as the '300dpi' specification comes from the typesetting department, who we need to follow as they are in charge of reproduction. Hopefully you have got the right versions already, or the person who has supplied the figures to you can let you have versions at 300dpi...many thanks in advance.

- Re: *percentage amount*. Under the crossheading 'Preparation of formulated product', you have made some adjustments, to create:

'We prepared 100 g of palm sap sugar by traditional production and cooked it in 1.0 litre of fresh water for fifteen minutes at 60° C to make an 11% palm sap sugar solution. Then, it was cooled in an open space for twenty minutes.

Furthermore, we also prepared 2.0 litres of mature coconut water (*Cocos nucifera* L.) and mixed it with the 1.0 litre of palm sap sugar solution (equivalent to 33% of palm sap sugar solution).'

Unfortunately, the detail about 33% is related to the earlier version of the text, before amounts and volumes were clarified. So it seems incorrect. If the palm sap sugar solution was 11%, then it seems that mixing a litre of it with 2 litres of another liquid (coconut water) will dilute the palm sap sugar content by 1/3rd; making 3 litres with 3.66% palm sap sugar solution per litre? NB, one potential solution here, rather than changing the percentage amount, would be to delete the phrase in brackets. What should be done please?

- Re: *small miscellaneous point*. In the paragraph after Figure 3, you adjusted the text to:

'GSI values in the KP3 diet were significantly ($F(3,8) = 10.492, P = 0.243$) higher than the KP1, KP2, and KP4 diets.' - In fact there was unfortunately a small error at the start of the sentence, so I just wanted to double-check it is correct like this (with the plural 'GSI values...').

- Re: *overlooked point*: In the names and addresses of institutions, there is sometimes 'West Sumatra' and sometimes 'Sumatera Barat' - and there isn't full consistency between the m/s and the manuscript system on this. Which should it be? Suggest make consistent to 'West Sumatera' in both places. Ok?

- Tables 1-4. Re: *the point about superscript letters*. Unfortunately the wording didnt quite work, for correct sense. Suggest use the following: 'Note: If the numbers in a row have different superscript letters, this indicates there is a significant difference between them ($P < 0.05$). If the numbers in a row have the same superscript letter, this means they show no significant difference ($P > 0.05$).'

[Quoted text hidden]

[Quoted text hidden]

[Quoted text hidden]

[Quoted text hidden]

[Quoted text hidden]

[Quoted text hidden]

[Quoted text hidden]

[Quoted text hidden]

[Quoted text hidden]

[Information Classification: General](#)



Webmail
Univ. Bung Hatta

azrita ubh <azrita31@bunghatta.ac.id>

Manuscript 124706 conditionally accepted for publication

azrita ubh <azrita31@bunghatta.ac.id>
To: "F1000.Research" <research@f1000.com>

Wed, Nov 16, 2022 at 9:32 PM

Dear

Jonathan

F1000 Research editorial team

Thank you for your email on November 16, 2022. We have revised manuscript No. 124706 (attached). At the same time, we also send all images that have been formatted with Jpeg 300 pi and list correction of the manuscript.

With best regards

Azrita

[Quoted text hidden]

5 attachments




Figure 3_pooled_TGC_001.jpg
492K





Figure 2_Polled_Growth coefficient__001.jpg
902K



-Fig 1. gut_001.jpg
1297K

 **124706 F1000 Research - Received JH2.doc**
3228K



List correction of manuscript No. 124706 .doc

25K

List correction of manuscript No. 124706 highlights magenta

1. All images have been formatted with Jpeg 300 pi
2. In Method (equivalent to 33% of palm sap sugar solution) has been deleted
3. In the paragraph after Figure 3, the text is revised to The GSI value of giant gourami was significantly different ($F_{(3,8)} = 10.492$, $P = 0.243$) between diets, and the GSI of giant gourami fed KP3 ration was higher than ration KP1, KP2, and KP4
4. Names and addresses of institutions have been revised and have complete consistency
5. Universitas Nahdlatul Ulama Sumatera Barat is the name of the institution
6. About superscript letters for Tables 1,2 and 3 have been adjusted based on suggestions of a reviewer

Corresponding author: Azrita

Effect of feed enriched by products formulated from coconut water, palm sap sugar, and mushroom on the chemical composition of feed and carcass, growth performance, body indices, and gut micromorphology of giant gourami, *Osphronemus goramy* (Lacepède, 1801), juveniles

Azrita undefined^{a*}, Hafrijal Syandri^b, Netti Aryani^c, Ainul Mardiah^d

^aDepartment of Biology, Faculty of Education, Universitas Bung Hatta, Padang, West Sumatera, 25133 Indonesia

^bDepartment of Aquaculture, Faculty of Fisheries and Marine Science, Universitas Bung Hatta, Padang, West Sumatera, 25133, Indonesia.

^cDepartment of Aquaculture, Faculty of Fisheries and Marine Science, Universitas Riau, Pekanbaru, 28293, Indonesia.

^dDepartment of Aquaculture, Faculty of Fisheries and Marine Science, Universitas Nahdlatul Ulama Sumatera Barat, Padang, West Sumatera, 28293, Indonesia.

Corresponding author: Azrita31@bunghatta.ac.id

ABSTRACT

Background: Giant gourami, *Osphronemus goramy* (Lacepede, 1801) is a freshwater species and Indonesia's most important commercial fish. Most giant gourami are produced by aquaculture. The first purpose of this study is to determine the effect of various newly formulated products on the amino acid composition of the diet and whole-body carcass. The second aim is to analyse the growth coefficient, body indices, and gut micromorphology.

Methods: A total of 1.0 litre of palm sap sugar solution and 2.0 litres of mature coconut water were formulated. 1 litre of product was added in turn to 2 g of *Aspergillus niger* (labeled CP2), 2 g of *Rhizopus oligosporus* (labeled CP3), and 2 g of *Saccharomyces cerevisiae* (labeled CP4), while, freshwater as a control (labeled CP1). Aquafeed was added to CP1, CP2, CP3, and CP4, to make diets labeled KP1, KP2, KP3, and KP4. The fish dosage was 150 ml/kg of feed. Juvenile giant gourami (initial weight 50±0.25 g and length 13.2±0.07 cm) were reared in triplicate net frames (2×1×1 m; water volume 1.5 m³/frame nets) in a freshwater concrete pond with a stocking density of 30 juveniles/net.

Results: The results supported our hypothesis that different product formulations have a significant effect ($P < 0.05$) on aquafeed nutrition and the whole-body carcass, growth coefficient, feed utilization, body indices, and gut micromorphology of giant gourami juveniles. The thermal growth coefficient strongly correlated with the daily growth coefficient ($r^2 = 91\%$). The KP3 diet contains a higher concentration of amino acids, which increased the growth coefficient, feed utilization, and carcass quality more than the other diets that we tested.

Conclusions: Diet KP3 contains higher total amino acids in diets and carcasses and leads to better growth for giant gourami.

Keywords: Giant gourami, amino acid profile, growth performance, feed efficiency, coconut water, gut micromorphology

Commented [HJ1]: Re: figures within the article.

#2 - Unfortunately, only one of the four attachments is at the resolution we need for reproduction (300 dpi); this is the one for Figure 3. Please therefore could you provide jpegs (or another suitable file format such as tiffs) at 300 dpi for the other figures to be used. Thanks.

Formatted: Highlight

Commented [HJ2]: On our system, the last part of the address that was entered currently reads: '...Universitas Bung Hatta, Padang, Sumatera Barat, 25113, Indonesia'.

#2 - Which address is correct please? They are different.

Formatted: Highlight

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Commented [HJ3]: #2 – you have made a change or two here. However the sentence starting 'Aquafeed was added...', which I queried, is still confusing, unfortunately. (It suggests that you add aquafeed to CP1 and then also add CP2, CP3, CP4 to the 'aquafeed plus CP1' mix.)

I think you must mean: 'Aquafeed was added to CP1, and then added to CP2, CP3, and CP4, to make diets labeled KP1, KP2, KP3, and KP4.' Please confirm.

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Introduction

In this decade, the production of capture fisheries has decreased; meanwhile, the demand for fish products for human consumption is increasing. Therefore, according to the Food and Agriculture Organisation, 60% of fisheries production in the future will come from aquaculture activities and this figure will continue to rise¹. The utilization of a variety of fish for aquaculture has now increased the need for commercial feed^{2,3,4,5}. At the same time, for aquaculture operations, the cost of aquafeed is still a significant challenge^{2,6,7,8}. On the other hand, commercial feed produced by factories still does not contain complete nutrition for fish growth, while being acknowledged for its positive effects on food safety^{9,10,11}. In this context, enriching fish feed with cost-effective natural ingredient resources is key to increasing feed nutrient quality and feed efficiency in commercial fish farming and ensuring the sustainability of aquaculture operations^{2,12,13}.

The target is fish feed that is wealthy in many important nutrients, including protein, fat, vitamins, and minerals that cultured fish can utilize to increase their growth rate and survival and that is beneficial for human health^{4,14,15,16}. Therefore, novel approaches have been developed by scientists to improve the nutrition of fish feeds, such as feed supplemented with EPA and DHA¹⁷, iodine and selenium¹⁰, methionine¹⁸, fish oil^{19,11}, and soybean oil²⁰. In addition, supplementing probiotics into the diet²¹ and supplemental glycine, prebiotics, and nucleotides in a soybean meal-based diet have been studied²².

The progress of aquaculture biotechnology has stimulated the interest of scientists in improving aquatic animal production, for example, to increase giant gourami production. One of the experimental techniques is to increase feed nutrition used for this purpose, such as, the use of fish meal and Azolla flour as a feed ingredient for giant gourami²³, and the utilization of new products formulated from water coconut, palm sap sugar, and fungus for the enrichment of commercial feed⁹. Additional research has involved a diet supplemented using glutamine²⁴, feed supplemented with a growth hormone²⁵, and substitute fish meal incorporating chicken feather²⁶. Whether using coconut water and palm sap sugar fermented with mushrooms affects the amino acid composition of the diet, body carcass, growth coefficient, and body indices is still not understood.

Coconut water has extraordinary nutritional value and contains supplements for health like minerals, amino acids, fatty acids, vitamins, enzymes, organic acids, and several phenolic compositions^{27,28,29,30}. Palm sap sugar also has health benefits due to its essential nutrient content, such as a low glycaemic index, and it contains antioxidants, vitamins, and minerals^{31,32,33,34}. Meanwhile, mushrooms have been widely used in fermentation due to their

Commented [HJ4]: #2 – It does not seem to work, putting 'FAO, 2018' in the text, as no other references like tis, which are given in the end-of-article References, are included within the text. So I have cut this and moved the spelt-out 'Food and Agriculture Organisation' within the sentence.
This seems to work well to me – if you agree?

Commented [SSR4]: We agreed

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

ability to degrade antigenic proteins in fish feed ingredients^{7,35,36}. Additionally, coconut water is a functional food that can protect the lens from diabetic cataract development in rats³⁷. Coconut water is also a treatment for burning pain during urination, dysuria, gastritis, increasing semen, and indigestion³⁸.

On the other hand, Azrita *et al.*⁹ have reported using new formulations of products containing coconut water and palm sap sugar that are fermented with various mushrooms involving a dosage of 300 ml/kg feed. Their newly formulated products can increase fatty acid levels in the diet and whole body carcasses. Besides that, they also improve giant gourami's growth performance and feed efficiency.

However, the effect of these new formulation products at a dosage of 150 ml/kg feed on the diet amino acid composition, and body meat's amino acid composition has not yet been analyzed. In line with that, the relationships between thermal growth coefficient and condition factor, daily growth coefficient, and feed utilization coefficient, including body indices parameters, as well as the gut micromorphology of giant gourami, have not yet been analyzed.

We hypothesized that commercial aquafeed combined with different newly formulated products at the dosage of 150 ml/kg feed could improve the amino acids compositions of the aquafeed and whole body carcass, body indices, and gut micromorphology. Hence, this investigation's first purpose was to analyze the effect of various newly formulated products on the diet's proximate compositions, amino acid composition, and whole-body carcass. The second aim was to analyze the impact of newly formulated products on the growth coefficient and relation to thermal growth coefficient, body indices, and gut micromorphology in giant gourami juveniles.

Methods

Ethical approval

The Research and Community Service Ethics Committee at Universitas Bung Hatta, West Sumatera, Indonesia approved this research (89/LPPM/Hatta/III-2022) which followed the ARRIVE guidelines. The Ministry of Education, Culture, Research and Technology of the Republic of Indonesia funded the research under grant No. 076/E5/PG.02.00 PT/2022 on March 16, 2022. Approval was given by the ethics committee to collect and rear juvenile gurami sago in the aquaculture laboratory, Faculty of Fisheries and Marine Science at Universitas Bung Hatta, Padang, West Sumatera, Indonesia. All efforts were made to relieve the suffering of experimental animals. Therefore, the animal did not suffer for this study, and

Commented [HJ6]: #2 - Please confirm this plural ('compositions') is correct

Formatted: Highlight

they were still in good condition when returned to the pond after research was completed. Where some fish were euthanized, this was carried out by piercing part of the fish's brain. Gurami sago fish are not classified as a protected animal according to Indonesian legislation.

Preparation of formulated product

We prepared 100 g of palm sap sugar by traditional production and cooked it in 1.0 litre of fresh water for fifteen minutes at 60° C to make an 11% palm sap sugar solution. Then, it was cooled in an open space for twenty minutes. Furthermore, we also prepared 2.0 litres of mature coconut water (*Cocos nucifera* L.) and mixed it with the 1.0 litre of palm sap sugar solution. The product was stored for ten minutes in a cool air-conditioned room. A total of 3.0 litres of the formulated product was divided into three parts of 1.0 litre each. We added 2 g of *Aspergillus niger* (labeled as CP2 product) to the first part of the formulated product solution, 2 g of *Rhizopus oligosporus* (labeled as CP3 product) to the second part, and 2 g of *Saccharomyces cerevisiae* (labeled as CP4 product) to the third portion. The CP2, CP3, and CP4 products were fermented for 48hrs in a jerry can (2.0 litres) using an Aerasi PUJIMAC, MAC-40 K 40 L/min. The products of CP2, CP3, and CP4 were used to enrich the nutrition of commercial aquafeed (781-2, PT. Japfa Comfeed Indonesia, Tbk) and labeled as the KP2, KP3, and KP4 diets. The aquafeed was supplemented with freshwater (labeled as the KP1 diet; placebo).

Preparation of experimental diets

Giant gourami juveniles were adapted for one month to standard feed, namely floating commercial aquafeed 781-2 (pellet size 2 mm), which contained 10.66% water content, 30.10% crude protein, 4.09% crude fat, 45.35% total carbohydrates, 2.5% ash, and 9.18% crude fibre. The minerals in the commercial feed were 280.08 mg/100 g Na, 1415.02 mg/100 g Ca, 1358.07 mg/100 g K, 1200.31 mg/100 g P, 292.03 mg/100 g Mg, 18.14 mg/100 g Fe, and 13.83 mg/100 g Zn. The aquafeed was added to freshwater to create the KP1 diet as observed, and the formulated CP2, CP3, and CP4 products were added to the aquafeed at a dosage of 150 ml/kg of feed to create the enriched fish diets. The formulated product added to the aquafeed was mixed manually with it for three minutes to obtain maximum homogenization and then the blend was dried in the open air for thirty minutes. Thereafter, it was given to the trial animal.

Experimental procedures and sampling

Formatted: Highlight

Formatted: Highlight

In the present study, we measured fish weight using AD-600i scales with 0.001 g accuracy (ACIS model number AD-600i, China). At the same time, a meter ruler with 1 mm accuracy was used to estimate the body length. A total of 360 sago strain juveniles of giant gourami were counted; the initial mean weight was 50 ± 0.25 g, and the initial length was 13.2 ± 0.07 cm. For rearing juveniles, twelve nets framed with 2 m^3 ($2 \times 1 \times 1$ m) PVC pipe (water volume of 1.5 m^3) were placed inside two freshwater concrete ponds with a size of 18 m^3 ($6 \times 2 \times 1.5$ m). This experiment consisted of four treatments and three replications, and each frame net was stocked with 30 juveniles. The giant gourami were fed the KP1, KP2, KP3, and KP4 diets three times a day (08:00, 12:00, and 17:00hrs) during the 90-day feeding trial. Juveniles of giant gourami were fed at a 3% body weight rate per day based on the percentage of stored biomass. Fish samples were collected every 30 days for body weight and length measurements. Ten fish per net frame were collected and anesthetized orally using clove oil. Then, their lengths and weights were measured. Prior to sampling, the fish fasted for 24hrs to empty their intestinal contents.

Proximate and amino acid composition

The diet samples and proximate carcass composition were analyzed using standard AOAC methods³⁹. The matter was dried to a constant weight at 105°C . We used the standard Kjeldahl method to analyse crude protein ($\text{N} \times 6.25$). We used the Soxhlet method with ether extraction to analyse crude lipids; the ash was incinerated at 550°C for 16 hrs, whereas gross energy was measured in a bomb calorimeter.- The amino acid composition was determined by using a high-performance liquid chromatography (HPLC) system consisting of a water 1525 binary HPLC pump, 717 autosamplers (water®), and water 2475 multi λ fluorescence detector optics (wavelengths: 250 nm for excitation and 395 nm for emission). It was hydrolysed in triplicate with 6 N hydrochloric acid for 24hrs at 11°C ⁴⁰.

Nutrient utilization and body indices

The growth coefficients in the fish experiments were measured by using the thermal growth coefficient (TGC), daily growth coefficient (DGC), total feed intake (FI), and protein efficiency ratio (PER) of giant gourami, assessed using the following formulae:

$$\text{TGC} = [(\text{final weight (g)})^{1/3} - (\text{initial weight (g)})^{1/3}] / (\text{mean water temperature } (^\circ\text{C})) \times \text{duration of rearing period (day)}] \times 1000$$

Commented [HJ7]: Sense of 'replicates' unclear...do you mean 'replications'?

#2 - I don't understand what 'replicate' means (or 'replicates'), I'm afraid. The word reads like an error. ('replicates' also reads like an error, it is incorrect usage.)

Do you mean you did the four treatments 3 times? (ie that you 'replicated' them 3 times? 'replicated' means 'repeated', as we know...)

I'd like to suggest that you phrase it a different way, so we can ensure the wording captures what you want...many thanks.

Formatted: Highlight

Formatted: Highlight

$$\text{DGC} = (\text{Wf}^{1/3} - \text{Wi}^{1/3}) / \text{duration of rearing period (day)} \times 100$$

$$\text{FI as feed (FI as feed in g/fish/day)} = \text{Total feed fed} / (n \times t)$$

$$\text{PER} = \text{wet weight gain} / \text{total protein intake}$$

Three fish from each net frame were sacrificed and dissected immediately to determine the Condition factor (CF), Viscerosomatic index (GSI%), Hepatosomatic index (HSI%), Visceral fat-somatic indexes (VFSl%), and Bilesomatic index (BSI) as given below:

$$\text{CF} = 100 \times [\text{weight of the juvenile (g)} / \text{Length of juvenile (cm}^3)]$$

$$\text{GSI} = 100 \times [\text{viscera weight (g)} / \text{whole body weight (g)}]$$

$$\text{HSI} = 100 \times [\text{liver weight (g)} / \text{whole body weight (g)}]$$

$$\text{VFSl} = 100 \times [\text{visceral fat weight (g)} / \text{whole body weight (g)}]$$

$$\text{BSI} = 100 \times [\text{Bile weight (g)} / \text{weight of liver (g)}]$$

Histological examination of the gut

For histological analyses, each gut specimen of the animal was cut into the foregut, midgut, and hindgut. Moreover, the cells were cleaned in saline solution and fixed in Bouin's fixative solution for 24hrs. After sequential dehydration steps in alcohol, the gut samples were embedded in paraffin. The implanted tissue blocks were sectioned at 5 μm , and sections were consistently stained with Haematoxylin-eosin and observed under a light microscope (Olympus IX71) equipped with Image-Pro Plus 7.0 software. The digitalized analysis measures the micrometer length of various enteric structures of gut images. We determined the average fold height (hF), fold width (wF), and enterocyte height (hMV) of the gut per slice (5 fields per individual sample) according to procedures described by Li *et al.*¹⁸. The specific measurement method of gut samples is shown in [Figure 1](#).




Figure 1. Transversal section photomicrographs of giant gourami juvenile foregut. (A) Fold height and fold width were analyzed in a lower magnification of objective lens of microscope (magnification $\times 100$), (B) Enterocytes height and microvilli height were analyzed using a higher magnification of an objective lens microscope (magnification $\times 200$). hF = fold height, wF = fold width, hE = enterocyte height, hMV = microvillus height (hematoxylin and eosin).

Pond water quality

The water quality values of the freshwater concrete ponds that were used to rear the giant gourami juveniles were recorded weekly. The water samples were collected at 10:00am at a depth of 20 cm from each concrete pond to determine the water temperature, dissolved oxygen, and pH value. In addition, we also measured the total alkalinity, hardness, and nitrates of the water in the pond experiments. A thermometer (Celsius scale) was used to measure water temperature. To measure water dissolved oxygen (O_2 ; $mg L^{-1}$), we used an oxygen meter (YSI Model 52, Yellow Instrument Co, Yellow Spring, OH USA). A digital pH meter (Mini 0–14 pH IQ, Scientific Cemo Science, Thailand) was used to determine the pH values of water in the experiments. The level of nitrate-nitrogen (NO_3-N ; $mg L^{-1}$), alkalinity ($mg L^{-1}$), and hardness ($mg L^{-1}$) were measured according to standard procedures⁴¹.

Calculations and statistical method

The data from this study were reported in the form of the mean \pm standard deviation for each treatment. Data were analysed using the SPSS 16.0 software package (SPSS; Chicago, IL). Normality was tested using the Kolmogorov–Smirnov statistic. Homogeneity was checked using absolute residuals according to Levine's test. One-way ANOVA was used to determine the treatment effect, followed by a post-hoc Duncan's multiple range test⁴². To create the figures, Microsoft Office Professional Plus 2019 was used.

Results

Proximate and amino acid profiles of the diets

Commercial feed supplemented with different formulated products with the dosage of 150 ml/kg of feed significantly affects the proximate composition of diets. One-way ANOVA results showed a marginal interaction among treatments in the case of protein content ($F_{(3,8)} = 1.522, P = 0.282$), fat ($F_{(3,8)} = 5.663, P = 0.022$), carbohydrates ($F_{(3,8)} = 1.862, P = 0.214$), crude fibre ($F_{(3,8)} = 1.445, P = 0.300$), and ash ($F_{(3,8)} = 0.272, P = 0.844$), and the total energy content ($F_{(3,8)} = 1.112, P = 0.400$) differed considerably ($P < 0.05$) among the four diets (Table 1). Duncan's Post-hoc test revealed that the protein content ($21.6967 \pm 0.17\%$) was significantly higher ($P < 0.05$) in the KP3 diet than in the other treatments, while the carbohydrate ($31.19 \pm 0.38\%$), crude fibre ($2.82 \pm 0.06\%$), and ash ($6.67 \pm 0.06\%$) contents were significantly higher ($P < 0.05$) in the KP3 diet than in the other diets. Conversely, the total energy content was 240.88 ± 0.74 (kg calories/100 g), which was significantly higher ($P < 0.05$) in the KP3 diets than in the KP1, KP2, and KP4 diets (Table 1).

The levels of free amino acids in the diets supplemented with different formulated products with a dosage of 150 ml/kg of feed are presented in Table 1. All types of amino acids in the diets of KP1, KP2, KP3, and KP4 were significantly different ($P < 0.05$), except for tryptophan, and there was no significant difference ($P > 0.05$) between KP2, KP3, and KP4. Among the essential amino acids, leucine and arginine were found in the highest amounts in the KP1, KP2, KP3, and KP4 diets. There was no significant difference ($P > 0.05$) in the alanine content between KP2 and KP3 diets and the cystine level in KP1 and KP3 diets. Of the nonessential amino acids, glutamic and aspartic acid represented a significant portion of all four diets.

The present study found significant differences in the overall free essential and nonessential amino acid pools in the KP1, KP2, KP3, and KP4 diets (Table 1). One-way ANOVA results exhibited a marginally significant interaction between experimental diets in terms of essential amino acids ($F_{(3,8)} = 11.371, P = 0.003$), nonessential amino acids ($F_{(3,8)} = 0.407, P = 0.752$), and overall amino acid pools (essential plus nonessential) ($F_{(3,8)} = 7.355, P = 0.011$). Duncan's Post-hoc test revealed that the free essential amino acids ($9.10 \pm 0.011\%$), nonessential amino acids ($12.91 \pm 0.004\%$), and overall amino acid pools (22.02%) were significantly higher ($P < 0.05$) in feed supplemented with CP3 products, followed by CP2, CP4, and CP1 products (Table 1).

Formatted: Highlight

Table 1. The experimental diets' proximate and amino acid composition (% dry matter). Mean \pm SD*. Note: If the numbers in a row have different superscript letters, this indicates there is a significant difference between them ($P < 0.05$). If the numbers in a row have the same superscript letter, this means they show no significant difference ($P > 0.05$).

	KP1	KP2	KP3	KP4
<i>Proximate composition</i>				
	%, dry weight basis			
Dry matter	38.42 \pm 0.25 ^a	38.27 \pm 0.01 ^a	37.59 \pm 0.16 ^a	38.41 \pm 0.10 ^a
Crude protein	19.68 \pm 0.41 ^a	20.27 \pm 0.13 ^b	21.70 \pm 0.18 ^c	20.44 \pm 0.10 ^b
Crude lipid	3.41 \pm 0.02 ^a	3.67 \pm 0.13 ^b	3.50 \pm 0.02 ^{ac}	3.48 \pm 0.04 ^{ab}
Carbohydrate	26.37 \pm 0.17 ^a	29.50 \pm 0.54 ^b	31.19 \pm 0.38 ^c	30.57 \pm 0.06 ^b
Crude fibre	2.23 \pm 0.05 ^a	2.36 \pm 0.01 ^b	2.82 \pm 0.06 ^c	2.45 \pm 0.06 ^b
Ash	2.75 \pm 0.03 ^a	6.66 \pm 0.05 ^b	6.57 \pm 0.04 ^c	6.67 \pm 0.06 ^b
Energy total (kg calorie/100 g)	240.87 \pm 0.38 ^a	234.41 \pm 0.30 ^b	240.88 \pm 0.74 ^{ac}	237.11 \pm 0.43 ^b
<i>Amino acid composition</i>				
EAA				
Leucine	1.36 \pm 0.01 ^a	1.42 \pm 0.01 ^b	1.46 \pm 0.01 ^c	1.36 \pm 0.01 ^d
Isoleucine	0.76 \pm 0.01 ^a	0.79 \pm 0.01 ^b	0.81 \pm 0.01 ^c	0.76 \pm 0.01 ^d
Lysine	0.95 \pm 0.01 ^a	1.10 \pm 0.01 ^b	0.98 \pm 0.01 ^c	1.20 \pm 0.01 ^d
Valine	0.86 \pm 0.01 ^a	0.94 \pm 0.01 ^b	0.96 \pm 0.01 ^c	0.89 \pm 0.01 ^d
Threonine	0.79 \pm 0.02 ^a	0.92 \pm 0.01 ^b	1.04 \pm 0.01 ^c	0.83 \pm 0.01 ^d
Arginine	1.02 \pm 0.01 ^a	1.19 \pm 0.01 ^b	1.30 \pm 0.01 ^c	1.03 \pm 0.01 ^d
Phenylalanine	0.67 \pm 0.01 ^a	0.93 \pm 0.01 ^b	1.05 \pm 0.01 ^c	0.77 \pm 0.01 ^d
Tyrosine	0.43 \pm 0.01 ^a	0.53 \pm 0.00 ^b	0.57 \pm 0.06 ^c	0.45 \pm 0.01 ^d
Methionine	0.18 \pm 0.01 ^a	0.26 \pm 0.01 ^b	0.30 \pm 0.01 ^c	0.21 \pm 0.01 ^d
Histidine	0.40 \pm 0.01 ^a	0.50 \pm 0.01 ^b	0.57 \pm 0.01 ^c	0.43 \pm 0.01 ^d
Tryptophan	0.06 \pm 0.01 ^a	0.11 \pm 0.01 ^b	0.07 \pm 0.00 ^{bc}	0.09 \pm 0.01 ^{bd}
NEAA				
Alanine	0.85 \pm 0.01 ^a	0.94 \pm 0.01 ^b	0.87 \pm 0.06 ^c	0.97 \pm 0.01 ^{bd}
Serine	1.01 \pm 0.01 ^a	1.12 \pm 0.01 ^b	1.23 \pm 0.01 ^c	1.01 \pm 0.01 ^d
Glycine	1.15 \pm 0.01 ^a	1.32 \pm 0.01 ^b	1.29 \pm 0.01 ^c	1.19 \pm 0.01 ^d

Formatted: Highlight

Formatted: Highlight

Formatted: Superscript

Formatted: Highlight

Formatted: Superscript

Formatted: Highlight

Formatted: Superscript

Formatted: Highlight

Formatted: Superscript

Commented [HJ8]: Query – there is 'ac' and 'ad' in superscript in this line.

This sort of thing occurs occasionally elsewhere. Is it just a case of typos please, or is it deliberate? (If it is erroneous, please go through each table and delete the unwanted letters.)

#2 – Actually, there appears to be no explanation of what any of the superscript letters mean. Please add a note at the bottom of the table about this, therefore – and also, for any other table where this applies...

Proline	1.01 ± 0.01 ^a	1.05 ± 0.01 ^b	1.03 ± 0.01 ^c	1.03 ± 0.02 ^d
Aspartic acid	1.25 ± 0.01 ^a	1.50 ± 0.01 ^b	1.40 ± 0.01 ^c	1.56 ± 0.01 ^d
Glutamic	2.15 ± 0.03 ^a	2.88 ± 0.03 ^b	2.59 ± 0.01 ^c	3.01 ± 0.03 ^d
Cystine	0.09 ± 0.01 ^a	0.07 ± 0.01 ^b	0.04 ± 0.01 ^c	0.09 ± 0.01 ^{ad}
∑EAA	7.56 ± 0.003 ^a	8.70 ± 0.003 ^b	9.03 ± 0.003 ^c	8.04 ± 0.003 ^d
∑NEAA	7.51 ± 0.008 ^a	8.88 ± 0.007 ^b	8.88 ± 0.004 ^c	8.84 ± 0.008 ^d
∑AA	15.07 ± 0.004 ^a	17.58 ± 0.002 ^b	17.91 ± 0.00 ^c	16.88 ± 0.003 ^d

* Values represent the means of triplicate samples.

Proximate and amino acid profile of the whole body of giant gourami

Commercial feed combined with a new formulation product significantly affected the proximate carcass composition of juvenile giant gourami. One-way ANOVA results showed a marginal interaction among group treatments in the case of protein contents ($F_{(3,8)} = 1.522$, $P = 0.282$), fat ($F_{(3,8)} = 5.663$, $P = 0.022$), carbohydrates ($F_{(3,8)} = 1.862$, $P = 0.214$), and crude fibre ($F_{(3,8)} = 1.445$, $P = 0.300$). Duncan's Post-hoc test revealed that the protein content ($21.69 \pm 0.17\%$), fat ($3.47 \pm 0.03\%$), carbohydrates ($31.18 \pm 0.37\%$), and crude fibre ($2.81 \pm 0.05\%$) were significantly higher ($P < 0.05$) in the KP3 diet than in the other treatments. Meanwhile, the carcass protein content of fish fed KP1, KP2, and KP4 was not significantly different ($P > 0.05$) between treatments. For the energy total, KP3 was significantly higher ($P < 0.05$) than the other treatments (Table 2). However, the moisture content of the carcass did not show any significant variation among the KP1, KP2, KP3, and KP4 diets.

The mean quantities of total amino acids in the carcasses of *O. goramy* fed different diets are given in Table 2. Lysine and leucine represented a significant portion of the essential amino acids of the whole body carcass, and methionine was present in small quantities in all of the whole-body meat. Of the nonessential amino acids, glutamic acid, aspartic acid, and alanine were the highest, and cystine was the lowest for all whole-body carcasses of giant gourami fed different diets. The levels of glutamic acid were significantly higher in carcasses of fish fed the KP3 diet than in those provided the KP1, KP2, and KP4 diets.

Table 2. Whole-body proximate and amino acid composition of giant gourami after a 90-day feeding trial. **Note:** If the numbers in a row have different superscript letters, this indicates

Formatted: Highlight

there is a significant difference between them ($P < 0.05$). If the numbers in a row have the same superscript letter, this means they show no significant difference ($P > 0.05$).

	KP1	KP2	KP3	KP4		
<i>Proximate composition</i>		%, dry wet basis				
Dry matter	64.59 ± 0.16 ^c	64.51 ± 0.34 ^c	64.14 ± 0.33 ^c	64.24 ± 0.12 ^c	Formatted: Highlight	
Crude protein	28.64 ± 0.28 ^a	28.07 ± 0.79 ^{ab}	28.85 ± 0.45 ^c	28.66 ± 0.44 ^{ad}	Formatted: Superscript	
Crude fat	2.79 ± 0.03 ^a	2.88 ± 0.02 ^b	2.67 ± 0.04 ^c	3.00 ± 0.02 ^d	Formatted: Highlight	
Carbohydrate	1.38 ± 0.01 ^a	1.99 ± 0.06 ^b	1.97 ± 0.09 ^c	1.31 ± 0.02 ^d	Formatted: Highlight	
Crude fibre	0.97 ± 0.02 ^a	0.68 ± 0.01 ^b	0.83 ± 0.02 ^c	0.95 ± 0.04 ^d	Formatted: Superscript	
Ash	1.63 ± 0.02 ^a	1.70 ± 0.02 ^b	1.54 ± 0.01 ^c	2.11 ± 0.04 ^d	Formatted: Highlight	
Energy total (kg calorie/100 g)	144.77 ± 1.58 ^a	155.48 ± 1.26 ^b	157.90 ± 0.91 ^c	149.60 ± 0.29 ^d	Formatted: Superscript	
<i>Amino acid composition</i>						
<i>EAA</i>						
Leucine	2.13 ± 0.01 ^a	2.37 ± 0.01 ^b	2.42 ± 0.01 ^c	2.26 ± 0.01 ^d		
Isoleucine	1.13 ± 0.01 ^a	1.25 ± 0.01 ^b	1.38 ± 0.01 ^c	1.19 ± 0.01 ^d		
Lysine	2.77 ± 0.01 ^a	3.16 ± 0.02 ^b	3.88 ± 0.01 ^c	2.86 ± 0.01 ^d		
Valine	1.26 ± 0.01 ^a	1.40 ± 0.01 ^b	1.32 ± 0.01 ^c	1.35 ± 0.01 ^d		
Threonine	1.38 ± 0.02 ^a	1.49 ± 0.01 ^b	1.43 ± 0.01 ^d	1.48 ± 0.01 ^d		
Arginine	1.58 ± 0.01 ^a	1.71 ± 0.01 ^b	1.63 ± 0.01 ^c	1.70 ± 0.01 ^d		
Phenylalanine	1.02 ± 0.01 ^a	1.11 ± 0.01 ^b	1.08 ± 0.01 ^c	1.11 ± 0.01 ^d		
Tyrosine	0.80 ± 0.01 ^a	0.84 ± 0.00 ^b	0.83 ± 0.01 ^c	0.85 ± 0.06 ^d		
Methionine	0.15 ± 0.01 ^a	0.21 ± 0.01 ^b	0.18 ± 0.01 ^c	0.16 ± 0.01 ^d		
Histidine	0.55 ± 0.01 ^a	0.56 ± 0.01 ^{ab}	0.59 ± 0.01 ^c	0.57 ± 0.01 ^d	Formatted: Highlight	
Tryptophan	0.08 ± 0.01 ^a	1.02 ± 0.01 ^b	1.08 ± 0.01 ^c	0.06 ± 0.00 ^d	Formatted: Highlight	
<i>NEAA</i>						
Alanine	1.86 ± 0.01 ^a	2.08 ± 0.01 ^b	2.92 ± 0.01 ^c	1.97 ± 0.01 ^d		
Serine	1.28 ± 0.01 ^a	1.31 ± 0.01 ^b	1.26 ± 0.01 ^c	1.31 ± 0.01 ^d	Formatted: Highlight	
Glycine	1.58 ± 0.01 ^a	1.68 ± 0.01 ^b	1.61 ± 0.01 ^c	1.77 ± 0.01 ^d		

Formatted: Highlight

Formatted: Superscript

Formatted: Highlight

Formatted: Superscript

Formatted: Highlight

Formatted: Superscript

Formatted: Superscript, Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Proline	1.06 ± 0.01 ^a	1.16 ± 0.01 ^b	1.08 ± 0.01 ^c	1.16 ± 0.01 ^d
Aspartic acid	2.71 ± 0.01 ^a	3.08 ± 0.01 ^b	3.79 ± 0.01 ^c	2.77 ± 0.01 ^d
Glutamic	4.36 ± 0.03 ^a	4.92 ± 0.01 ^b	4.97 ± 0.01 ^c	4.66 ± 0.01 ^d
Cystine	0.06 ± 0.01 ^a	0.09 ± 0.01 ^b	0.06 ± 0.01 ^c	0.05 ± 0.01 ^d
∑EAA	12.68 ± 0.003 ^a	15.13 ± 0.005 ^b	15.82 ± 0.001 ^c	13.61 ± 0.008 ^d
∑NEAA	12.91 ± 0.007 ^a	14.32 ± 0.01 ^b	15.69 ± 0.002 ^c	13.50 ± 0.001 ^d
∑AA	25.59 ± 0.003 ^a	29.45 ± 0.04 ^b	31.51 ± 0.001 ^c	27.11 ± 0.004 ^d

* Values represent the means of triplicate samples.

When the overall quantities of total essential and nonessential amino acids were compared, the whole-body carcass amino acid content was significantly lower ($P < 0.05$) in fish fed the KP1 diet than in those fed the KP2, KP3, and KP4 diets (Table 2). The number of amino acids (essential plus nonessential) in the carcasses of fish fed the KP3 diet was significantly higher than that in fish fed the KP1, KP2, and KP4 diets.

Growth coefficient and survival

The growth coefficient and feed utilization of the giant gourami juveniles displayed significant differences among the diets. One-way ANOVA results exhibited a marginally significant difference between experimental diets in the case of the thermal unit growth coefficient ($F_{(3,8)} = 153.99$, $P = 0.458$), and daily growth coefficient ($F_{(3,8)} = 59.88$, $P = 0.288$), while total feed intake (% BW day⁻¹) ($F_{(3,8)} = 14.938$, $P = 0.56$), and protein efficiency ratio ($F_{(3,8)} = 15.78$, $P = 0.29$) also showed significant differences ($P < 0.05$) among the treatment diets (Figure 2).




Figure 2. Growth coefficient and feed utilization of the giant gourami juveniles reared under different diets during 90 days of the experiment period. (A) Thermal growth coefficient (TGC), (B) daily growth coefficient (DGC), (C) feed intake (FI), and (D) protein efficiency ratio (PER). The mean value and standard deviation (mean \pm SD) are presented for giant gourami ($n = 3$). Different superscripts in the bar diagram of the giant gourami juvenile TGC, DGC, FI, and PER indicate significant differences among other diets ($P < 0.05$, One-way ANOVA Duncan Post-Hoc)

Furthermore, the thermal growth coefficient (TGC) has often been used to predict growth performance and production performance of aquaculture using water temperature at the fish-

rearing location. This study presents the relationship between thermal growth coefficient and condition factor, daily growth coefficient, and protein efficiency ratio (Figure 3). The thermal growth coefficient had strong relationships with the condition factor ($r^2 = 0.777$, figure 3A), daily growth coefficient ($r^2 = 0.999$, figure 3B), and protein efficiency ratio ($r^2 = 0.749$, figure 3D), while the thermal growth coefficient had a moderate relationship with the feed intake ($r^2 = 0.699$, figure 3C).




Figure 3. Relationships between thermal growth coefficient and condition factor (A), daily growth coefficient (B), feed intake (C) and protein efficiency ratio (D) for giant gourami (*O. gourami*) over 90 days.

Condition factor and body indices of giant gourami after 90 days of feeding

The condition factor was significantly different between diets ($F_{(3,8)} = 19.98, P = 0.566$) in the present study; while the GSI, HIS, and VFSI displayed marginally significant differences between diets. The HIS was significantly ($F_{(3,8)} = 5.389, P = 0.500$) higher in the KP3 diet, but KP1, KP2, and KP4 diets had no significant differences among them (Table 3). The GSI value of gourami was significantly different ($F_{(3,8)} = 10.492, P = 0.243$) between diets, and the GSI of giant gourami fed KP3 ration was higher than ration KP1, KP2, and KP4. The VFSI was not considerably different among the KP1, KP2, and KP4 diets. The Duncan's post-hoc test revealed that the HIS ($1.30 \pm 0.13\%$), GSI ($4.15 \pm 0.36\%$), and VFSI ($2.75 \pm 0.34\%$) were significantly higher ($P < 0.05$) in the KP3 diet than in the other diets. Meanwhile, BSI showed no significant difference ($P > 0.05$) among the treatment diets (Table 3).

Formatted: Highlight

Table 3. Mean (\pm SD) value condition factor and body indices of giant gourami during the 90-day experimental period. Note: If the numbers in a row have different superscript letters, this indicates there is a significant difference between them ($P < 0.05$). If the numbers in a row have the same superscript letter, this means they show no significant difference ($P > 0.05$).

Formatted: Highlight

Growth coefficients	KP1	KP2	KP3	KP4
Condition factor (CF)	2.45 ± 0.09^a	2.90 ± 0.07^b	2.92 ± 0.13^c	2.61 ± 0.04^d
Viscerosomatic index (GSI%)	3.20 ± 0.21^a	3.77 ± 0.09^b	4.15 ± 0.36^c	3.17 ± 0.02^d
Hepatosomatic (HIS%)	0.97 ± 0.05^a	1.06 ± 0.19^{ab}	1.30 ± 0.13^c	1.04 ± 0.12^{ad}
Visceral fat-somatic indexes (VFSI%)	2.15 ± 0.13^a	2.29 ± 0.22^{ab}	2.75 ± 0.34^c	1.74 ± 0.21^{ad}
Bilesomatic (BSI%)	10.11 ± 0.76	10.58 ± 1.01	10.48 ± 1.28	10.29 ± 0.77

Gut micromorphology

The gut morphometric measurements of giant gourami juveniles are presented in Table 4. Fish gut micromorphology was significantly affected by different feeds. One-way ANOVA results showed a significant effect of feed differences between groups in terms of foregut fold height ($F_{(3,8)} = 816.70, P = 0.135$), foregut fold width ($F_{(3,8)} = 129.34, P = 0.974$), height of the foregut ($F_{(3,8)} = 169.80, P = 0.882$), and microvillus height of the foregut ($F_{(3,8)} = 56.01, P = 0.285$). The Duncan's post-hoc test demonstrated that the foregut fold height ($434.13 \pm$

1.76 μm), fold width ($53.23 \pm 0.88 \mu\text{m}$), enterocyte height ($27.42 \pm 0.42 \mu\text{m}$), and microvillus height ($2.79 \pm 0.45 \mu\text{m}$) were significantly higher ($P < 0.05$) in fish fed the KP3 diet than those fed the other diets. For the midgut, one-way ANOVA results showed a significant interaction among treatments in the case of fold height ($F_{(3,8)} = 5602.628$, $P = 0.055$), fold width ($F_{(3,8)} = 129.341$, $P = 0.974$), enterocyte height ($F_{(3,8)} = 169.809$, $P = 0.882$), and microvillus height ($F_{(3,8)} = 56.016$, $P = 0.285$). The Duncan's post-hoc test showed that the fold height of the midgut ($324.96 \pm 1.43 \mu\text{m}$), fold width ($61.50 \pm 1.02 \mu\text{m}$), and enterocytes ($32.82 \pm 0.54 \mu\text{m}$) were significantly higher ($P < 0.05$) in fish fed the KP3 diet, whereas microvillus height was significantly higher in fish fed the KP2 diet (Table 4). Fish fed the KP3 diet showed a higher fold height of the hindgut ($F_{(3,8)} = 5459.01$, $P = 0.066$), fold width ($F_{(3,8)} = 271.94$, $P = 0.865$), enterocyte height ($F_{(3,8)} = 299.180$, $P = 0.821$), and microvillus height ($F_{(3,8)} = 253.57$, $P = 0.316$).

Pond water quality

The pond water quality values of the giant gourami juvenile rearing freshwater concrete pond were recorded; water temperatures, dissolved oxygen (DO), total alkalinity, hardness, pH, and nitrates were in the range of typical values as given by WHO/FAO, as shown in Table 5.

Table 4. Gut micromorphology of giant gourami juveniles fed different diets for 90 days. Mean values with different superscript letters in the same line are significantly different ($P < 0.05$).

	Foregut				Midgut				Hindgut			
	hF (μm) ^a	wF (μm) ^b	hE (μm) ^c	hMV (μm) ^d	hF (μm)	wF (μm)	hE (μm)	hMV (μm)	hF (μm)	wF (μm)	hE (μm)	hMV (μm)
KP1	336.17±5.59 ^a	51.30±0.85 ^a	26.21±0.43 ^a	2.56±0.45 ^a	227.50±0.16 ^a	47.16±0.78 ^a	24.31±0.31 ^a	1.64±0.03 ^a	213.92±0.19 ^a	42.91±0.59 ^a	20.22±0.25 ^a	1.49±0.02 ^a
KP2	343.43±1.38 ^b	52.14±0.86 ^b	26.84±0.44 ^b	2.77±0.45 ^b	274.61±1.21 ^b	58.12±0.97 ^b	29.87±0.49 ^b	1.85±0.01 ^b	243.51±1.07 ^b	53.01±0.88 ^b	28.00±0.46 ^b	1.64±0.01 ^b
KP3	434.13±1.76 ^c	53.2±0.88 ^a	27.42±0.42 ^c	2.79±0.45 ^c	324.96±1.43 ^c	61.50±1.02 ^c	32.82±0.54 ^c	1.80±0.03 ^c	305.60±1.35 ^c	60.02±0.99 ^c	29.54±0.49 ^c	1.77±0.02 ^c
KP4	321.18±1.42 ^d	50.20±0.83 ^b	25.62±0.79 ^d	2.31±0.07 ^d	228.45±1.01 ^b	56.95±0.95 ^d	29.19±0.48 ^d	1.69±0.01 ^d	217.69±0.96 ^d	61.64±1.03 ^d	24.32±24.32 ^d	1.40±0.01 ^d

^a hF = fold height
^b wF = fold width
^c hE = enterocyte height
^d hMV = microvillus height

Formatted: Highlight

Formatted: Highlight

Table 5. The average values and range of water quality parameters in the concrete pond during the 90-days of experiment.

Water quality parameters	n	Mean ± SD	Range	WHO/FAO limits	References
Water temperatures ($^{\circ}\text{C}$)	45	28.01 ± 1.06	27 - 30	25-33	Prokoso <i>et al.</i> ⁴³
Dissolved oxygen (mg/L)	14	6.01 ± 0.14	5.80 - 6.20	3-5	Syandri <i>et al.</i> ⁴⁴
Total alkalinity (mg/L as CaCO_3)	14	58.09 ± 3.33	52.5 - 62.5	120	Boyd <i>et al.</i> ⁴⁵
Hardness (mg/L as CaCO_3)	14	66.34 ± 1.32	65 - 68.5	168	Boyd <i>et al.</i> ⁴⁵
pH	14	7.48 ± 0.19	7.2 - 7.8	6.5 - 9.0	Boyd <i>et al.</i> ⁴⁵
Nitrates (mg/L)	14	0.04 ± 0.01	0.03 - 0.05	0.2 - 219	Boyd and Tucker ⁴⁶

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Discussion

The chemical analysis of fish feed is essential because it provides valuable information to aquafeed nutritionists concerned with readily available sources of proximate and amino acid compositions, including minerals and vitamins. This study investigated the nutritional quality of fish feed enriched with three different formulation products and one as a placebo. Dietary protein levels for giant gourami ranged from 19.68 to 21.70%. Overall, the crude protein content in the feed of this study was within the ranges observed by other authors^{47,48,49}. The giant gourami belongs to the **trophic level** of herbivorous fish⁵⁰. Generally, herbivorous fish require a lower dietary protein level than carnivorous fish^{51,49}. Reducing the protein content of aquafeed is one method to increase continuous fish farming, by diminishing feed costs and reducing the impact on the aquatic environment^{2,52}. The fat content of the feed ranged from 3.41 to 3.67%, which is similar to the feed fat content for juvenile grass carp, *Ctenopharyngodon idella*⁵³, and lower than the feed fat content for the herbivorous fish *Ancistrus cirrhosis*⁴⁸ and for rearing rohu, *Labeo rohita*⁵⁴. At the same time, the carbohydrate content of all feed treatments ranged from 26.37 to 31.19%, and the energy total (kg calorie/100 g) was between 234.41 and 240.87. Although protein content as an energy source for the maintenance and growth of giant gourami is relatively low, energy can be acquired from either protein or nonprotein sources, i.e., fat and carbohydrates.

In the present study, the commercial fish feed was enriched with natural sources, i.e., formulated products of mature coconut water and palm sap sugar fermented with various fungi (*Aspergillus niger*, *Rhizopus oligosporus*, and *Saccharomyces cerevisiae*). In the recent past, the dose used was 300 ml/kg of feed. This method is a new approach that has been developed by Azrita *et al.*⁹ to improve feed nutrition and whole-body carcasses, covering fatty acids, the atherogenic index and thrombogenic, feed efficiency, and growth performance of giant gourami. Here, we continued the investigation by reducing the feed dose to 150 ml/kg. This study's results found that supplementing feed with newly formulated products can increase feed nutrition, covering amino acids in diet and body meat, and the growth coefficient of giant gourami. Several authors have reported increasing feed nutrition and maximizing the digestive enzyme activity of aquacultured fish by providing feed supplemented with EPA and DHA¹⁷, iodine and selenium¹⁰, methionine¹², fish oil^{19, 11}, and soybean oil²⁰. In addition, the provision of feed has been supplemented with probiotics²¹, glycine, and prebiotics²². In this study, mature coconut water and palm sap sugar solution fermented with various fungi were used to supplement fish feed. In addition to coconut water

and palm sugar, mushrooms also play a role in increasing feed nutrition. However, it's better to use *Rhizopus oligosporus*. As in the present study, Varzakas⁵⁵ and Vong *et al.*⁵⁶ showed that *Rhizopus oligosporus* can produce various extracellular enzymes. *Aspergillus niger* has a high capacity to degrade antigenic proteins, including carbohydrases, proteases, lipases, and phosphatases, when used for fermenting plant-sourced fish feed ingredients^{12,57}. *Saccharomyces cerevisiae* is one of the most acclaimed microorganisms. Its effectiveness is due to its useful composition, such as "β-glucans, nucleic acids, mannan oligosaccharides and chitin," which are used for fermented ingredients^{7,58}.

The amino acid composition can be used to assess feed quality. Leucine, arginine, and glutamic acid were the most abundant free amino acids in the KP1, KP2, KP3, and KP4 diets.

Similarly, in other studies on fish feed, such as feed for largemouth bass, *Micropterus salmoides*, the feeds were supplemented with glycine, prebiotics, and nucleotides in a soybean meal-based diet²². Feed for pacu, *Piaractus mesopotamicus*, was supplemented with an essential amino acid⁵⁹, and feed for snubnose pompano, *Trachinotus blochii*, was supplemented with different levels of protein⁶⁰. Apparently, supplementing feed with different ingredients is common, and in other species, leucine, arginine, and glutamic acid were the most abundant FAAs. Conversely, methionine levels were low in all experimental feeds. Methionine is one amino acid that must be available in fish feed because methionine is needed to protect body cells from stress. For optimal growth of juvenile hybrid grouper, 1.89% methionine is required in the feed¹⁸. The experimental feed contained 0.18–0.30% methionine, but whether this amount is sufficient for the needs of giant gourami is poorly understood.

In the current study, the nonessential amino acid compositions were slightly higher than the essential amino acid compositions in all the experimental diets. It was higher in the KP3 diet than the other diets. In contrast, the essential amino acids of fish feed for snubnose pompano were slightly higher than the nonessential amino acids content⁶⁰. This difference may be caused by differences between freshwater fish and marine fish. As in the present study, Prabu *et al.*⁶⁰ reported that different dietary protein levels also caused different pools of FAAs, including limiting essential amino acid types in the diet⁵⁹ and supplemental glycine, prebiotic, and nucleotide levels in the soybean meal-based diet²². In the present study, this difference in FAA content is caused by various mushrooms used in the formulated products.

Giant gourami juveniles fed the KP3 diet showed higher levels of glutamic acid, aspartic acid, leucine, and lysine and lower levels of tyrosine, methionine, histidine, tryptophan, and

Formatted: Highlight

Commented [HJ11]: #2 – it reads like there is a word missing before 'feed supplemented' if none is added

cystine in their carcasses than those fed other diets. The carcasses of giant gourami fed the KP3 diet showed the highest sum of FAAs compared to cultured fish fed the KP1, KP2, and KP4. The differences in the FAA profile in the whole-body carcasses of giant gourami could be related to the fungus type used in the formulated products for enriched feed. Each type of mushroom has a different function depending on the fermented fish feed ingredients and is correlated with the whole-body carcass amino acids^{12,57}. The FAA profile differences could be related to different aspects, such as diet composition⁶¹, dietary protein level⁶², and methionine levels in the diet¹⁸, including the water quality of the ponds⁶³. This study does not analyse the relationship between growth performance and FAA profile or pond water quality. Several authors have reported that the physiological parameters of water quality and animal body composition are usually interrelated^{64,60}. The present study did not examine whether the difference in FAAs in the whole-body carcass is correlated to pond water quality.

The lower weight gain of fish fed the KP1 diet compared to fish fed the KP2, KP3, and KP4 diets shows that a deficiency of either fungus in the formulated product for the enriched diet could lower the protein content and related sum amino acids, leading to the inhibition of giant gourami growth. In addition, it also affects feed intake and feed conversion ratios. The low protein efficiency ratio and daily growth coefficient in fish provided the insufficient KP1 diet were perhaps due to an amino acid imbalance. The amino acid content of the KP2, KP3, and KP4 diets increased, ranging from 16.88% to 17.91% after fermentation. The increase may be due in part to the increased protein content in the KP2, KP3, and KP4 diets, which was in line with the results of Jannatullah *et al.*⁵⁷ and Li *et al.*¹², who found that *Aspergillus niger* and *Aspergillus awamori* fermentation increased the amino acid content of soybean meal by 2.56% and 15.56%, respectively. In addition, Dawood *et al.*³⁶ stated that the essential amino acid profile was changed after fermentation by *Saccharomyces cerevisiae*. This might result from the different fungi used having different utilization patterns for amino acids in this study. It influences the growth performance and nutrient utilization of giant gourami juveniles. We found that the methionine proportion was lower in the diets in the current study. In addition, methionine is an essential amino acid that plays a unique role in protein structure and metabolism¹⁸. It is possible that *Aspergillus niger*, *Rhizopus oligosporus*, and *Saccharomyces cerevisiae* fermentation promoted the conversion of specific amino acids to methionine. However, the exact mechanisms need to be studied further.

In the present study, the thermal growth coefficient (TGC) strongly correlated with the daily growth coefficient (DGC). Because faster daily fish growth requires a quality diet and

constant water temperature during the rearing period, in this study, water temperature ranged from 27 to 30°C, and dissolved oxygen was between 5.8 and 6.2 mg /L. According to Besson *et al.*⁶⁵, higher daily energy availability in the diet can lead to faster-growing fish, which is supported by constant water temperature and higher daily oxygen levels. The thermal growth coefficient had an essential change in environmental value⁶⁶. Therefore, it was very important to keep the water temperature and dissolved oxygen constant in the aquaculture locations. At the same time, 78% of TGC values were determined by the condition factor connected to whole body weight and the total fish length. TGC of Atlantic cod, *Gadus morhua*, is influenced by body size and condition factors⁶⁷.

In this study, a higher value of TGC was detected in fish fed KP3; the effect is that the daily growth coefficient, and the protein efficiency ratio is better. Conversely, decreasing TGC has two effects, *i.e.*, a **slow fish** growth and lowered daily feed intake. Many scientists state that in aquaculture operations, net yield (kg/m³) depends upon TGC fluctuation, feed intake, and daily oxygen consumption^{65,68,69}.

In the present study, feed enrichment with different formulated products did not affect HIS or VFSI except in the KP3 diet. Whereas GSI is influenced by differences in diet, it did not affect BSI. The condition factor of largemouth bass, *Micropterus salmoides* (1.49–1.52%), fed enriched 1–2% EPA + DHA¹⁷ was different from the value (0.68) reported by Arriaga-Hernandez *et al.*⁷⁰ for white snook (*Centropomus viridis*) juveniles fed a 15% replacement of fish meal with soybean meal. Moreover, Hassan *et al.*⁷¹ reported condition factor values ranging from 1.52 to 2.95 and an HSI between 1.4 and 1.5 for *Lates calcarifer* under different feeding rates (3–9% body weight d⁻¹). Barbosa *et al.*⁷² reported VSI and LSI values of 2.24 and 3.86, respectively, for *Centropomus parallelus* fed a commercial diet. On the other hand, Syed *et al.*⁶⁴ also reported HSI and VSI values of 3.41 and 4.90, respectively, for *Oreochromis niloticus* at different levels of aloe vera extract as feed additives. In our study, the VSI of *O. goramy* ranged from 3.17 to 4.15, and the LSIs were between 1.74 and 2.75, both higher than those recorded at different stocking densities of *O. goramy*⁴⁴. The high content of visceral fat observed in fish fed the KP3 diet might be explained by the diet having fat contents that exceed the needs of giant gourami juveniles and by the reduced energy expenditure of fish that are confined to rearing frame nets. Therefore, further analysis is necessary to determine the optimum dosage of the formulated product for the enrichment of feed to improve the growth performance of *O. goramy*.

For fish, the gut plays a significant role in absorbing nutrients, which is closely related to feed utilization^{18,73}. Rossi *et al.*²² demonstrated that the development of enterocytes affected the nutrient-absorbing efficiency of the gut of *Micropterus salmoides*. Feeding *Lates calcarifer* juveniles with the same basal diet supplemented with 1% probiotic yeast, *Saccharomyces cerevisiae*, and lactic acid bacteria, *Lactobacillus casei*, revealed a higher number of gut mucosal goblet cells and increased microvillous length⁷⁴. In contrast, substituting as much as 12.5–25% soya protein concentrates with lupin (*Lupinus albus*) meal in carp (*Cyprinus carpio*) diets does not significantly affect the villi length and villi width of the gut⁷⁵. In the current study, enriched feed with products supplemented from coconut water, palm sap sugar, and fungus significantly affected the micromorphology and gut size. The fold height, fold width, enterocyte height, and microvilli of fish fed the KP3 diet were higher than those of fish fed the KP1, KP2, and KP4 diets. The KP3 diet is a relevant formulated product to enrich commercial feed to promote the development of the gut in animal experiments, which may somewhat describe the significant growth performance and feed efficiency used in this study.

Furthermore, the micromorphology gut size of giant gourami is smaller than that of juvenile hybrid grouper¹⁸, turbot, *Scophthalmus maximus*¹², largemouth bass, *Micropterus salmoides*²², and common carp, *Cyprinus carpio*⁷⁵. The trophic food habits of fish may also affect the gut's hF, wF, hE, and hMV size because these habits are correlated with the digestibility coefficient. Under natural conditions, giant gourami is an herbivorous fish, while grouper, largemouth bass, and turbot are predatory fish, and common carp are omnivorous. Whether giving fish from different trophic levels the same diet affects the size of gut hF, wF, hE, and hMV is poorly understood.

Conclusions

The present investigation observed that feed enriched with newly formulated products made from mature coconut water and palm sap sugar, and fermented with various mushrooms, given to fish in a dose of 150 ml/kg substantially affected the amino acid composition of the diet and whole-body carcass of giant gourami juveniles. It also affected the growth coefficient, feed utilization, body indices, and gut micromorphology size. The thermal growth coefficient had a strong relationship with the daily growth coefficient ($r^2 = 91\%$) and a moderate relationship with the feed intake ($r^2 = 69\%$). The CP3 formulation was optimal for feed quality, and the KP3 diet was optimal for body carcass, growth coefficient,

body indices, and the ability of the intestines for feed absorption. Thus, our study also informs fish farmers about culturing good quality giant gourami and fulfilling nutrition requirements for food security.

Data availability

Underlying data

Figshare: Underlying data for 'Effect of feed enriched by products formulated from coconut water, palm sap sugar, and mushroom on the chemical composition of feed and carcass, growth performance, body indices, and gut micromorphology of giant gourami, *Osphronemus goramy* (Lacepède, 1801), juveniles'.

<https://doi.org/10.6084/m9.figshare.20407647>⁷⁶

This project contains the following underlying data:

- Table 1. Raw data of the experimental diets' proximate composition
- Table 2. Raw data of amino acid of feed experimental
- Table 3. Raw data of whole body carcass proximate composition
- Table 4. Raw data of amino acid of whole-body carcass
- Table 5. Daily growth coefficient, feed utilization and body indices of giant gourami after 90 days of feeding.
- Table 6. Raw data gut micromorphology of giant gourami juveniles fed different diets for 90 days

Data are available under the terms of the [Creative Commons Attribution 4.0 International License](#) (CC-BY 4.0).

Competing interests

We, as the authors of this article, disclose no competing interests.

Grant information

The Ministry of Education, Culture, Research, and Technology of the Republic of Indonesia funded this research under grant number: 076/E5/PG.02.00. PT/2022.

Acknowledgments

We are grateful to the Ministry of Education, Culture, Research, and Technology of the Republic of Indonesia for funding this [research](#).

Commented [HJ12]: Please confirm you have checked that the body is happy for this acknowledgment to appear.

#2 – Suggest deleting the new line added, for convention, as it 'sounds odd'. The reason I asked if you had permission to thank the Ministry, is that F1000 always checks this; but it does not need to be stated or suggested in the Acknowledgements.

References

1. Food and Agriculture Organization (FAO): The state of world fisheries and aquaculture: Opportunities and challenges. Food and Agriculture Organization of the United Nations. 2018; p. 209.
2. Hua K, Cobcroft JM, Cole A, *et al.*: The future of aquatic protein: Implications for protein sources in aquaculture diets. *One Earth*. 2019; 1(3): 316–329.
3. Gómez B, Munekata PES, Zhu Z, *et al.*: Challenges and opportunities regarding the use of alternative protein sources: Aquaculture and Insects. *Adv. Food Nutr. Res.* 2019; 89: 259-295.
4. Yuru Z, Ronghua L, Chaobin Q, *et al.*: Precision nutritional regulation and aquaculture. *Aquac. Rep.* 2020; 18: 100496.
5. Nagappan S, Das P, AbdulQuadir M, *et al.*: Potential of microalgae as a sustainable feed ingredient for aquaculture. *J. Biotechnol.* 2021; 341: 1-20.
6. Hasimuna O, Maulu S, Monde C, *et al.*: Cage aquaculture production in Zambia: Assessment of opportunities and challenges on Lake Kariba, Siavonga district. *Egypt. J. Aquat. Res.* 2019; 45: 281-285.
7. Dawood MAO, Koshio S: Application of fermentation strategy in aquafeed for sustainable aquaculture. *Rev Aquac.* 2019; 1-16.
8. Junaidi, Syandri H, Azrita, *et al.*: Floating cage aquaculture production in Indonesia: Assessment of opportunities and challenges in Lake Maninjau. *AIMS Environ Sci.* 2022; 9(1): 1-15.
9. Azrita, Syandri H, Aryani N, *et al.*: The utilization of new products formulated from water coconut, palm sap sugar, and fungus to increase nutritional feed quality, feed efficiency, growth, and carcass of gurami sago, *Osphronemus goramy* (Lacepède, 1801) juvenile. *F1000Research*. 2021;10: 1121.
10. Barbosa V, Maulvault AL, Anacleto P, *et al.*: An enriched feed with iodine and selenium from natural and sustainable sources to modulate farmed gilthead seabream (*Sparus aurata*) and common carp (*Cyprinus carpio*) fillets elemental nutritional value. *Food Chem. Toxicol.* 2020;140: 111330.
11. Duarte FOS, Faula FG, Prado C.S, *et al.*: Better fatty acid profile in fillets of Nile tilapia (*Oreochromis niloticus*) supplemented fish oil. *Aquaculture*. 2020; 534: 736241.
12. Li X, Wu X, Dong Y, *et al.*: Effects of dietary levels on growth, feed utilization and related gene expression of juvenile hybrid grouper (*Epinephelus fuscoguttatus* ♀ × *Epinephelus lanceolatus* ♂). *Aquaculture*. 2019; 502: 153-161.
13. Nagarajan D, Varjani S, Lee D, *et al.*: Sustainable aquaculture and animal feed from microalgae: Nutritive value and techno-functional components. *Renew. Sust. Energ. Rev.* 2021;150: 111549.
14. Lall SP, Dumas A: Nutritional requirements of cultured fish. *Feed and Feeding Practices in Aquaculture*, 20115; 53-109.
15. Mohanty BP, Mahanty A, Ganguly S, *et al.*: Nutritional composition of food fishes and their importance and providing food and nutritional security. *Food Chem.* 2019; 293: 561-570.
16. Beltrán JMG, Esteban MA: Nature-identical compounds as feed additives in aquaculture. *Fish Shellfish Immunol.* 2022; 123: 409-416.
17. Yadav AK, Rossi W, Habte-Tsion, *et al.*: Impacts of dietary eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA) level and ratio on the growth, fatty acid

- composition, and hepatic-antioxidant status of largemouth bass (*Micropterus salmoides*):- *Aquaculture*. 2020; 59: 735683.
18. Li X, Mu W, Wu X, *et al.*: The optimum methionine requirement in diet of juvenile hybrid grouper (*Epinephelus fuscoguttatus* ♀ × *Epinephelus lanceolatus* ♂): Effects on survival, growth performance, gut micromorphology and immunity. *Aquaculture*. 2020; 520: 735014.
 19. Aryani N, Azrita, Mardiah A, *et al.*: Influence of feeding rate on the growth, feed efficiency, and carcass composition of the giant gourami (*Osphronemus goramy*). *Pakistan J. Zool.* 2017; 49(5): 1775-1781.
 20. Godoy AC, Oliveira SO, Oxford JH *et al.*: Soybean oil for Nile tilapia (*Oreochromis niloticus*) in finishing diets: Economic, zootechnical and nutritional meat improvements. *Aquaculture*. 2019; 734324.
 21. Ringø E: Probiotics in shellfish aquaculture. *Aquac. Fish.* 2020; 5(1): 1-27.
 22. Rossi W, Allen KM, Habte-Tsion H-M, *et al.*: Supplementation of glycine, prebiotic, and nucleotides in soybean meal-based diets for largemouth bass (*Micropterus salmoides*): Effects on production performance, whole-body nutrient composition and retention, and intestinal histopathology. *Aquaculture*. 2021; 532: 736031.
 23. Afriyanti EK, Hasan ODS, Djunaidi IS: Growth performance of giant gourami, *Osphronemus goramy* fed with a combination of fish meal and Azolla flour, Azolla microphyll. *Jurnal Iktiologi Indonesia*. 2020; 20(2): 133-14.
 24. Andriani J, Setiawati M, Sunarno MTJ: Diet digestibility and growth performance of giant gourami juvenile, *Osphronemus goramy* fed on a diet supplemented using glutamine. *Jurnal Iktiologi Indonesia*. 2019; 19(1): 1-11.
 25. Budi DS, Alimuddin, Suprayudi MA: Growth response and feed utilization of giant gourami (*Osphronemus goramy*) juvenile feeding different protein levels of the diets supplemented with recombinant growth hormone. *HAYATI J. Biosci.* 2015; 22 (1): 12-19.
 26. Wibowo WP, Sumidjan I, Rachmawati D.: Analysis of relative growth rate, feed efficiency and survival of giant gourami, *Osphronemus goramy* juvenile through substituting chicken feather flour silage in artificial feed. *Journal of Aquaculture Management and Technology*. 2017; 6(2): 51-58.
 27. Segura-Badilla O, Lazcano-Hernández M, Kammar-García A, *et al.*: Use of coconut water (*Cocos nucifera* L) for the development of a symbiotic functional drink. *Heliyon*. 2020; 6(3): e03653.
 28. Kumar M, Saini SS, Agrawal PK, *et al.*: Nutritional and metabolomics characterization of the coconut water at different nut developmental stages. *J. Food Compos. Anal.* 2021; 96: 103738.
 29. Paixão LB, Brandão GC, Araujo RGO, *et al.*: Assessment of cadmium and lead in commercial coconut milk employing HR-CS GF AAS. *Food Chem.* 2019; 284: 259-263.
 30. Lakshmanan J, Zang B, Wright K, *et al.*: Tender coconut water suppresses hepatic inflammation by activating AKT and JNK signaling pathways in an in vitro model of sepsis. *J. Funct. Foods*. 2020; 64: 103637.
 31. Ho CW, Wan Aida WM, Maskat, *et al.*: Changes in volatile compounds of palm sap (*Arenga pinnata*) during the heating process for production of palm sugar. *Food Chem.* 2007; 102: 1156-1162.
 32. Saputro AD, de Walle DV, Dewettinck K.: Physicochemical properties of coarse palm sup sugars as a natural alternative sweetener. *Food Biosci.* 2020; 38: 100780.
 33. Heryeni H: The virtues of palm sugar & product development strategy. Lampung Mangkurat University Press. (In Indonesian). 2016.

34. Sarma C, Mummaleti G, Sivanandham V, *et al.*: Anthology of palm sap: The global status, nutritional composition, health benefits & value added products. *Trends Food Sci. Technol.* 2022; 119: 530-549.
35. Garrido-Galand S, Asensio-Grau A, Calvo-Lerna J, *et al.*: The potential of fermentation on nutrition and technological improvement of cereal and legume flours: A review. *Int. Food Res. J.* 2021; 145: 110398.
36. Dawood MAO, Eweedah NM, Khalafalla MM, *et al.*: *Saccharomyces cerevisiae* increases the acceptability of Nile tilapia (*Oreochromis niloticus*) to date palm seed meal. *Aquac. Rep* 2020; 17: 100314.
37. Zhang X, Peng L, Dai Y, *et al.*: Anti-cataract effects of coconut water in vivo and in vitro. *Biomed. Pharmacother* 2021; 143: 112032.
38. Prades A, Dornier M, Diop N, *et al.*: Coconut water uses, composition and properties: a review. *Fruits.* 2012; 67(2): 87–107.
39. AOAC, Official Methods of Analysis. 15th Edn.: Association of Official Analytical Chemists (AOAC), Washington, DC., USA. 1990.
40. Cohen SA: Amino acid analysis using pre-column derivatization with 6-aminoquinolyl-N-hydroxysuccinimidyl carbamate. In: *Protein Sequencing Protocols*, Smith B.J. (Ed.). 2nd Edn. 2003; vol. 211, Humana Press Inc., Totowa, NJ., ISBN: 9781592593422, pp: 143-154.
41. APHA: Standard Methods for Examination of Water and Wastewater. 1995. 19th Edn. American Public Health Association, Washington DC, USA.
42. Duncan DB: Multiple ranges and multiple F tests. *Biometrics.* 1955; 11: 1-42.
43. Prokoso VA, Pouil S, Chahyanti W, *et al.*: Fluctuating temperature regime impairs growth in giant gourami (*Osphronemus goramy*) larvae. *Aquaculture.* 539: 736606.
44. Syandri H, Mardiah A, Azrita, *et al.*: Effect of stocking density on the performance of juvenile gurami sago (*Osphronemus goramy*) in the synthetic sheet pond. *Pak J Zool.* 2020; 52(2): 717-726.
45. Boyd CE, Tucker CS, Somridhivej B.: Alkalinity and hardness: Critical but elusive concepts in aquaculture. *J. World Aquac. Soc.* 2016; 47(1): 6-41.
46. Boyd CE, Tucker CS: Pond aquaculture water quality management. Kluwer Academic Publishers, Boston. 1988.
47. Timalsina P, Yadav CNR, Lamsal GP, *et al.*: Effect of stocking density and source of animal protein on growth and survival of rainbow trout fingerling in flow-through system at Nuwakot Nepal. *Aquac. Rep.* 2017; 8: 58-64.
48. Vucko MJ, Cole AJ, Moorhead J, *et al.*: The freshwater macroalga *Oedogonium intermedium* can meet the nutritional requirements of the herbivorous fish *Ancistrus cirrhosis*. *Algal Res.* 2017; 27: 21-31.
49. Yu H, Liang H, Ren M, *et al.*: A study to explore the effects of low dietary protein levels on the growth performance and nutritional metabolism of grass carp (*Ctenopharigodon idella*) fry. *Aquaculture.* 2022; 546: 737324.
50. Syandri H, Azrita, Sumiarsih E, *et al.*: Nutrient loading and farm characteristics of giant gourami fish aquaculture systems in Lake Maninjau, Indonesia: Basic knowledge of production performance. *F1000 Research.* 2021; 10: 378.
51. Omasaki SK, Janssen K, Besson M, *et al.*: Economic values of growth rate, feed intake, feed conversion ratio, mortality and uniformity for Nile tilapia. *Aquaculture.* 2017; 481: 124-132.
52. Arenas M, Alvarez-Gonzalez A, Barreto A, *et al.*: Effect of dietary carbohydrates on growth performance, feed efficiency and glucose metabolism in common snook

- (*Centropomus undecimalis*) and yellowtail snapper (*Ocyurus chrysurus*) juveniles. *Aquaculture*. 2021; 543: 736958.
53. Tian J, Li Y, Zhang K, *et al.*: Dietary Pennisetum sinense Roxb meal reduces lipid accumulation and alleviates enteritis in juvenile grass carp, *Ctenopharingodon Idella*. *Aquac. Rep.* 2021; 21: 1000915.
 54. Ali S, Kaviraj A: Aquatic weed *Ipomoea aquatica* as a feed ingredient for rearing Rohu, *Labeo rohita* (Hamilton). *Egypt. J. Aquat. Res.* 2018; 44(4): 321-325.
 55. Varzakas T: *Rhizopus oligosporus* mycelial penetration and enzyme diffusion in soya bean tempee. *Process Biochemistry*. 1998; 33(7): 741-747.
 56. Vong WC, Hua XY, Liu SQ: Solis-state fermentation with *Rhizopus oligosporus* and *Yarrowia lipolytia* improved nutritional and flavor properties of okara. *LWT*. 2018; 90: 316-322.
 57. Jannathulla R, Dayal JS, Ambasankar K, *et al.*: Effect of *Aspergillus niger* fermented soybean meal and sunflower oil cake on growth carcass composition and hemolymph indices in *Peneus vannamei* Boone, 1931. *Aquaculture*. 2018; 486: 1-8.
 58. Huang L, Ran CHS, Ren P, *et al.*: Effects of dietary *Saccharomyces cerevisiae* culture or live cells with *Bacillus amyloliquefaciens* spores on growth performance, gut mucosal morphology, hsp70 gene expression, and disease resistance of juvenile common carp (*Cyprinus carpio*). *Aquaculture*. 2015; 438: 33-38.
 59. Oliveira TS, Khan KU, Boarartt, AZ, *et al.*: Evaluation of the optimum dietary essential amino acid pattern for adult pacu (*Piarachus mesopotamicus*). *Aquaculture*. 2021; 540: 736686.
 60. Prabu DL, Ebeneezar S, Chandrasekar S, *et al.*: Influence of graded level of dietary protein with equated level of limiting amino acids on growth, feed utilization, body indices and nutritive profile of snubnose pompano, *Trachinotus blochii* (Lacepede, 1801) reared in low saline water. *Anim. Feed Sci. Technol.* 2020; 269: 114685.
 61. Hua K, Bureau D: Estimating changes in essential amino acid requirements of rainbow trout and Atlantic salmon as a function of body weight or diet composition using a novel factorial requirement model. *Aquaculture*. 2019; 513: 734440.
 62. Lu X, Peng D, Chen X *et al.*: Effect of dietary protein levels on growth, muscle composition, digestive enzymes activities, hemolymph biochemical indices and ovary development of pre-adult red swamp crayfish (*Procombarus clarkia*). *Aqua Rep.* 2020; 100542.
 63. Fatima S, Komal W, Manzoor F, *et al.*: Analysis of the growth performance, stress, profile of fatty acids and amino acids and cortisol in *Tilapia (Oreochromis niloticus)*, cultured at high stocking density using in-pond raceway system. *Saudi J. Biol. Sci.* 2021; 28(12): 7422-7431.
 64. Syed R, Masood Z, Hasan HU, *et al.*: Growth performance, hematological assessment and chemical composition of Nile tilapia, *Oreochromis niloticus* (Linnaeus, 1758) fed different levels of Aloe vera extract as feed additives in a closed aquaculture system. *Saudi J. Biol. Sci.* 2022; 29: 296-303.
 65. Besson M, Vandeputte M, van Arendonk JAM, *et al.*: Influence of water temperature on the economic of growth in fish farming: The case of sea bass (*Dicentrarchus labrax*) cage farming in the Mediterranean. *Aquaculture*. 2016; 462: 47-55.
 66. Besson M, Aubin J, Komen H, *et al.*: Environmental impacts of genetic improvement of growth rate and feed conversion ratio in fish farming under rearing density and nitrogen output limitations. *J. Clean. Prod.* 2015; 116(10): 100-109.
 67. Jobling M: The thermal growth coefficient (TGC) model of fish growth: A cautionary note. Short Communication. *Aquac. Res.* 2003; 34: 581-584.

68. Zhou X, Wang J, Huang L, *et al.*: Modelling and controlling dissolved oxygen in recirculating aquaculture systems based on mechanism analysis and an adaptive PID controller. *Comput Electron Agric.* 2022; 192: 106583.
69. Sun M, Hassan SG, Li D: Models for estimating feed intake in aquaculture: A review. *Comput. Electron. Agric.* 2016; 127: 425-428. Publisher Full Text.
70. Arriaga-Hernandez D, Hernandez C, Martinez-Montano E, *et al.*: Fish meal replacement by soybean products in aquaculture feeds for white snook, *Centropomus viridis*: Effect on growth, diet digestibility, and digestive capacity. *Aquaculture.* 2021; 530: 735823.
71. Hassan HU, Ali QM, Ahmad N, *et al.*: Assessment of growth characteristics, the survival rate and body composition of Asian sea *Lates calcarifer* (Bloch, 1790) under different feeding rates in closed aquaculture system. *Saudi J. Biol. Sci.* 2021; 28: 1324-1330.
72. Barbosa MC, Jatoba A, Vieira FN, *et al.*: Cultivation of Juvenile Fat Snook (*Centropomus parallelus*, Poey, 1860) Fed Probiotic in Laboratory Conditions. *Braz Arch Biol Technol.* 2011; 54, (4): 795-801.
73. Mohammadi M, Imani A, Farhangi M, *et al.*: Replacement of fishmeal with processed canola meal in diets for juvenile Nile tilapia (*Oreochromis niloticus*): Growth performance, mucosal innate immunity, hepatic oxidative status, liver and intestine histology. *Aquaculture.* 2019; 518: 734824.
74. Siddik MAB, Foysal, MJ, Fotedar R, *et al.*: Probiotic yeast *Saccharomyces cerevisiae* coupled with *Lactobacillus casei* modulates physiological performance and promotes gut microbiota in juvenile barramundi, *Lates calcarifer*. *Aquaculture.* 2022; 546: 737346.
75. Anwar A, Wan AH, Omar S, *et al.*: The potential of a solid-state fermentation supplement to augment white lupin (*Lupinus albus*) meal incorporation in diets for farmed common carp (*Cyprinus carpio*). *Aqua Rep.* 2020; 17: 100348.