

Analysis of Fatty Acids and Amino Acids of Three Local Freshwater Bagridae Fish Species In The Kampar Kanan River Indonesia For Food Security

by Azrita Azrita

Submission date: 12-Jan-2024 09:47AM (UTC+0700)

Submission ID: 2269713314

File name: 2024_Analysis_of_Fatty_Acids_and_Amino_Acids_of_Three_Local.pdf (3.55M)

Word count: 6294

Character count: 30957

Research Article

Analysis of Fatty Acids and Amino Acids of Three Local Freshwater Bagridae Fish Species in the Kampar Kanan River, Indonesia, for Food Security

Azrita Azrita ¹, Hafrijal Syandri ¹, HazlinaAhamad Zakeri ²,
Harfiandri Damanhuri ¹, and Netti Aryani ³

¹Faculty of Fisheries and Marine Science, Bung Hatta University, 25131 Padang, West Sumatra Province, Indonesia



²Faculty of Science and Marine Environment, Universiti Malaysia Terengganu, Malaysia

³Faculty of Fisheries and Marine Science, Riau University, 28293 Pekanbaru, Riau Province, Indonesia

Correspondence should be addressed to Hafrijal Syandri; syandri_1960@bunghatta.ac.id

Received 13 July 2023; Revised 15 December 2023; Accepted 19 December 2023; Published 4 January 2024

Academic Editor: Farid Mansouri

Copyright © 2024 Azrita Azrita ³³ et al. ⁴⁴ This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Fish have become an irreplaceable dietary source of animal protein, especially among households with low socioeconomic status in rural and urban areas of Indonesia. This study is aimed at analysing the proximate composition, minerals, fatty acids, and amino acids of three local Bagridae fish species in the Kampar Kanan river, Indonesia. The standard AOAC method was employed to examine the proximate composition of the carcass, and the analysis of amino acids and fatty acids was conducted through HPLC and GC-MS, respectively. The mineral content was determined using AAS. The nutrient composition results of *Hemibagrus nemurus*, *Hemibagrus wyckii*, and *Mystus nigriceps* revealed that the protein content was 24.26%, 22.57%, and 21.39% (% dry weight), whereas the total lipid content was 6.64%, 7.47%, and 7.75%, respectively. Regarding mineral contents, the calcium levels ranged from 1.49 to 1.66 mg/g, iron levels from 28.35 to 40.36 µg/g, and zinc levels from 24.03 to 54.46 µg/g. Among the fatty acids, palmitic acid was the most abundant in all three species, accounting for 25.59–30.70% of the total fatty acids. Additionally, significant amounts of C18:1 (oleic acid), C18:0 (stearic acid), and C20:4 (arachidonic acid) were also detected as primary fatty acids. The calculated atherogenic index values in the three species of Bagridae fish ranged from 0.73 to 0.99, while the thrombogenic index values varied between 0.54 and 0.75. The predominant amino acids found in the three species of Bagridae fish were glutamic acid with their concentrations ranging from 9.10 to 24.34%. These results indicate that consuming the meat of these three freshwater Bagridae fish species caught in the wild does not pose any health risks to consumers. They can be considered a safe and suitable food source with good nutritional quality.

1. Introduction

Fish serve as a high-quality protein source that is essential for satisfying dietary protein requirements, especially in areas where other protein sources are few or difficult to obtain [1, 2]. Their accessibility can aid in ensuring a steady supply of food, particularly for communities whose main source of income is fishing or aquaculture [3]. Fish can contribute to a more varied selection of protein sources in communities, resulting in a diet that is more nutritious and balanced. Dietary fish can treat nutrient deficits and enhance general nutritional health. Fish are generally considered safe

and offer a lower risk of contamination with harmful substances compared to some land-based animal products. A reliable indicator of a fish's quality, nutritional value, physiological status, and environment is its chemical makeup, which is found in fish flesh [4].

Hemibagrus nemurus, locally known as “baung,” *Hemibagrus wyckii*, known as “geso,” and *Mystus nigriceps*, known as “ingir-ingir,” are three species of catfish found in the Kampar Kanan River, Indonesia [5, 6]. Due to their high market value, these species have been identified as significant economic resources for rural communities [7]. Therefore, the analysis of the nutritional quality of fish meat in

the study area is very important to determine whether it is beneficial to human health.

Knowing the composition of fatty acids and amino acids in fish meat is an essential factor that should not be ignored [8]. This information provides valuable insight into the nutritional quality of fish and the value of more nutritious and healthy foods [9]. In addition, a better understanding of the composition of fatty acids and amino acids can assist in selecting a suitable feed [10], developing a balanced diet [11], and maintaining optimal fish meat nutrition [12].

Omega-3 polyunsaturated fatty acids (PUFAs), particularly docosahexaenoic acid (DHA) and eicosapentaenoic acid (EPA) derived from fish oil, exhibit anti-inflammatory and antitumorigenic cancer properties, making them crucial as immune-boosting nutrients [13, 14]. Studies have demonstrated that ω -3 and ω -6 PUFAs exhibit beneficial effects in the management of cardiovascular diseases and cancers [15, 16]. The fatty acid composition of PUFAs can differ among different types of fish, including both freshwater and marine species [17, 18]. Amino acids (AAs), such as cysteine, arginine, tyrosine, glycine, proline, and serine, play a vital role in illness and stress situations and in the prevention of inflammation and repair processes in fish intestines [19, 20].

Furthermore, knowledge of the atherogenicity index (AI) and thrombogenicity index (TI) of fatty acids in the three local freshwater fish species of Bagridae can contribute to improving dietary guidelines. The AI and TI can be used to develop recommendations aimed at improving heart health [21, 22]. These recommendations may involve selecting fish with favourable fatty acid profiles, allowing individuals to choose the most suitable fish species based on their specific needs [17, 23]. Therefore, this study is aimed at examining the nutritional composition profile, mineral content, amino acids, fatty acids, and AI and TI of three freshwater fish species, namely, *H. nemurus*, *H. wyckii*, and *M. nigriceps*. These three fish species are generally consumed by people who live around the Kampar Kanan River in Kampar Regency, Riau Province, Indonesia.

2. Material and Methods

2.1. Animal Material. In this study, a total of 27 Bagridae fish belonging to three distinct species (*H. nemurus*, *H. wyckii*, and *M. nigriceps*) were examined. The specimens were obtained from local fishers operating in the upper stretches of the Kampar Kanan River in Kampar Regency, Riau Province (Figure 1). Sampling took place between June and August 2022 and was verified by the Fisheries Department, Faculty of Fisheries and Marine Science, Bung Hatta University, Indonesia. The samples were placed in plastic bags and transported to the laboratory in an insulated icebox. The samples were then grouped based on their respective sampling locations.

2.2. Biometry Measurement. Upon reaching the laboratory within 10 hours, the fish samples were subjected to gutting and washing. Each specimen was weighed (TW) and measured to determine the standard length (SL) as well as the

depth or maximum height (H). TW was measured using a balance scale (OHAUS Model CT 6000 USA) with a precision of 0.01 g. The length measurement was taken from the tip of the mouth to the end of the upper lobe of the caudal fin, representing the total body length, using a metre ruler with an accuracy of 1 mm. The height measurement involved a vertical assessment of the body's maximum height measured using a Digital Sekhmet Sigma Vernier calliper with an accuracy of 1 mm. The condition factor (CF) was calculated using the formula $CF = (TW/SL^3) \times 100$.

2.3. Proximate, Amino Acid, and Mineral Content Analysis. The proximate carcass composition was assessed following the AOAC standard methods [24]. The samples were dried to a constant weight at 105°C. The crude protein content was analysed using the standard Kjeldahl method, which involves multiplying the nitrogen content by a factor of 6.25. The Soxhlet method with ether extraction was employed to analyse crude lipids. The ash content was obtained by incinerating the samples at 550°C for 16 h. The amino acid composition was determined using a high-performance liquid chromatography (HPLC) system, which consisted of a Waters 525 binary HPLC pump, Waters 717 autosamplers, and Waters 2475 multi- λ fluorescence detector (with excitation at 250 nm and emission at 395 nm). The samples were hydrolysed in triplicate with 6 N hydrochloric acid for 24 h at 110°C [25].

For mineral data composition (Na, Mg, Ca, K, P, Fe, Cu, Mn, and Zn), the ashed feed sample and carcass were dissolved in 1 ml of hydrochloric acid (35% v/v Suprapur®, Merck) and then filtered with cellulose filter paper (Watchman No. 1, International Ltd; Maidstone, UK) and diluted to an appropriate concentration for each elemental mineral. P levels were analysed with a Perkin-Elmer AA spectrophotometer mod 3110 (Norwalk, CT, USA).

2.4. Fatty Acid Analysis. The fish meat was examined utilizing the gas chromatography-mass spectrometry (GC-MS) method. The method of Folch et al. (1957) modified by Rajion et al. [26] was employed to perform the total fat extraction. This involved using a solvent system consisting of chloroform and methanol at a 2:1 (v/v) ratio. The process of transmethylation was conducted using a solution of 14% methanolic boron trifluoride. The fatty acid composition of the meat was then analysed at the SIG Laboratory, Accredited Testing Laboratory-LP-184-IDN.

The nutritional quality of lipids (AI and TI) was calculated using the following equations [27]:

$$AI = \frac{[(C12:0 + 4 \times C14:0 + C16:0)]}{[(\sum MUFA + \sum n-6) + \sum n-3]}$$

$$TI = \frac{[(C14:0 + C16:0)]}{[(0.5 \times \sum MUFA) + (0.5 \times \sum n-6) + (3 \times \sum n-3) + (\sum n-3)/\sum n-6]}$$
(1)

where AI is the atherogenic index, TI is the thrombogenic index, C12:0 is the lauric acid, C14:0 is the myristic acid, C16:0 is the palmitic acid, $\sum MUFA$ is the sum of the concentrations of all monosaturated fatty acids, $\sum n-6$ is the sum of the concentrations of n-6 polyunsaturated fatty acids,




FIGURE 1: Map of Kampar Regency, Riau Province, and sampling locations of the three species of Bagridae fish.

and $\sum n - 3$ is the sum of the concentrations of n-3 polyunsaturated fatty acids.

16
 2.5. Data Analysis. Data analysis was performed using the SPSS 16.0 software package (SPSS; Chicago, IL). The homogeneity of variances was assessed using Levene's test.

One-way ANOVA was carried out to determine the parameters of proximate and amino acid composition, including the composition of fatty acids and 55 nutritional quality of lipids (AI and TI) for each species, followed by the post hoc Duncan's multiple range test [28]. The results are reported as the mean values \pm standard errors for each parameter.

3. Results

Table 1 presents the average wet weight, standard length, height, and the results of meat nutritional composition analysis for three indigenous freshwater Bagridae species in the Kampar Kanan river, Indonesian. The moisture content (% wet weight) ranged from 82.40 to 85.39%. Among the species examined, *M. nigriceps* exhibited the lowest protein content at 21.39%, while *H. nemurus* displayed the highest value at 24.26%. However, *M. nigriceps* had a higher mineral content, including iron and zinc.

Table 2 summarizes the fatty acid composition (% total fatty acid) for the three Bagridae fish species. C16:0 was the predominant fatty acid in the three species of Bagridae fish, with percentages ranging from 25.09 to 30.71%. The additional prominent fatty acids found were C18: 1, C18:0, and C22:6. *H. nemurus* exhibited a ω -6/ ω -3 ratio below 1 (0.82), while *H. wyckii* and *M. nigriceps* had ratios of 1.11 and 1.02, respectively. The obtained PUFA/saturated fatty acid (SFA) (P/S) ratio varied between 0.61 and 0.76. The AI values varied between 0.76 and 0.99 and the TI between 0.40 and 0.63 (Table 3).

Table 4 presents the amino acid composition (% of total protein) of the three Bagridae species. Glutamic acid was the dominant amino acid, ranging from 19.29 to 24.34%, followed by aspartic acid, which ranged from 9.21 to 11.27%. The lysine content was consistent, ranging from 9.67 to 9.87%. In *H. nemurus*, the levels of various amino acids ranged from 0.86 to 24.35%. Similarly, in *H. wyckii*, the amino acid levels ranged from 0.76 to 21.58%, while in *M. nigriceps*, the levels ranged from 0.74 to 19.29%.

4. Discussion

The results of this study provide important new information about the nutritional makeup of the studied species of Bagridae fish. The biometric measures of the fish yielded important information about their physical traits and health. Researchers are able to determine the average weight of fish at various lengths by examining the link between length and weight for a particular species. This knowledge is essential for understanding the trends in fish population growth, identifying potential shifts in growth rates over time, and assessing the general health and welfare of fish [29]. A variety of factors, including sex, age, maturity level, size, stomach fullness level, sampling strategies, sample sizes, and environmental circumstances, have an impact on fish health and the parameters that determine length-weight connections in fish [30]. However, none of these factors were considered in our study. In general, according to the theory put forth by Bagenal and Tesch [31], heavier fish of a given length are in better physiological condition. The condition factor is also used in fishery science to compare the health, fatness, and condition of fish [32]. A condition factor of >1 is favourable, indicating a good feed nutrition level and suitable living habitat [33]. Thus, based on the findings in this study, the Kampar Kanan River provides a suitable habitat for the fish, ensuring their accessibility and availability for community consumption.

The fish samples exhibited good nutritional profiles, with high amounts of proteins and lipids as well as moisture content contributing to their overall nutritional value, according to the results of proximate analysis and mineral composition (see Table 1). These results imply that the fish species studied provide a possible source of nutrients to solve issues with food security. The study of the mineral composition revealed the existence of critical minerals such as phosphorus (P), calcium (Ca), magnesium (Mg), iron (Fe), and zinc (Zn) in the fish samples in addition to proteins and lipids. All three Bagridae fish significantly differed from each other in terms of their crude ash content and mineral profile (see Table 1). However, in general, P and Ca are the two highest minerals found in all fish. Both of these macrominerals are good for human bone and tooth health, as key components of hydroxyapatite, the mineral matrix that provides strength and structure to bones and teeth [34].

In the current study, the meat of three species of Bagridae fish from the Kampar Kanan River in Indonesia was found to contain an abundance of microminerals such as Fe and Zn. *M. nigriceps* has been reported to exhibit higher mineral content, including Fe and Zn. This species is categorized as a small indigenous fish (SIF) with a maximum standard length of approximately 10.29 cm. According to [35], SIFs are generally known to be rich in Ca, Fe, and Mn. On the other hand, scale flour from three species of freshwater fish, such as *Osphronemus goramy*, *Cyprinus carpio*, and *Oreochromis niloticus*, also contains high levels of macrominerals and microminerals [36].

In this study, all three Bagridae fish had a total fat content of approximately 7–8% by weight, with a higher water content ranging from 82.40 to 85.39% (Table 1). These values are similar to those found in wild and cultured sea bass *Dicentrarchus labrax* [37]. Additionally, the fish species can be grouped into different categories, including high-fat fish (>8%), medium-fat fish (4–8%), low-fat fish (2–4%), and lean fish (<2%) [38]. According to the classification established by [38], the three indigenous Bagridae fish species are characterized as having a moderate fat content level, which is reflected by the high levels of DHA and EPA in the meat of *H. nemurus*, *H. wyckii*, and *M. nigriceps* (Table 2). Therefore, we recommend that consuming the meat of these three fish species can indirectly boost food security in rural and urban communities.

All three Bagridae fish species analysed in the study contained arachidonic acid (C20:4), at 12.13% for *H. nemurus*, 12.49% for *H. wyckii*, and 12.41% for *M. nigriceps*. Arachidonic acid serves as a precursor for synthesizing prostaglandins and leukotrienes [39]. Leukotrienes are crucial in allergic responses, inducing smooth muscle contraction and increasing vascular permeability [40]. Additionally, all three species also contained EPA ranging from 1.35 to 5.24% and DHA ranging from 11.56 to 15.65%. The increased consumption of omega-3 fatty acids, such as EPA and DHA, has been linked to a decreased risk of pain development in older individuals and to the prevention of Alzheimer's disease [41, 42].

As determined by the current study, the ω -6/ ω -3 ratios of the three Bagridae fish species ranged from 1:0.8 to

TABLE 1: Results of biometric, proximate, and mineral composition of three species of Bagridae fish.

	Species		
	<i>Hemibagrusnemurus</i>	<i>Hemibagruswyckii</i>	<i>Mystusnigriceps</i>
<i>Biometric measurements</i>			
Total weight (g)	389.99 ± 24.96 ^a	1390.33 ± 168.29 ^b	17.57 ± 1.53 ^c
Standard length (cm)	28.16 ± 0.53 ^a	44.14 ± 1.98 ^b	10.29 ± 0.15 ^c
Height (cm)	8.44 ± 0.16 ^a	8.82 ± 0.39 ^b	2.57 ± 0.03 ^c
Condition factor	1.63 ± 0.08	1.58 ± 0.07	1.56 ± 0.08
<i>Proximate composition</i>			
Crude protein (% DW)	24.26 ± 0.87 ^a	22.57 ± 0.37 ^b	21.39 ± 0.15 ^c
Crude fat (% DW)	6.64 ± 0.03 ^a	7.47 ± 0.02 ^b	7.75 ± 0.40 ^c
Crude ash (% DW)	1.94 ± 0.02 ^a	2.30 ± 0.09 ^b	1.59 ± 0.02 ^c
Moisture (% WW)	82.40 ± 2.51	85.39 ± 2.36	83.75 ± 2.21
<i>Mineral composition</i>			
<i>Macrominerals (mg/g)</i>			
Sodium	0.99 ± 0.00 ^a	1.61 ± 0.01 ^b	1.61 ± 0.02 ^c
Magnesium	1.12 ± 0.01 ^a	0.57 ± 0.02 ^b	1.15 ± 0.01 ^c
Calcium	1.66 ± 0.00 ^a	1.55 ± 0.02 ^b	1.49 ± 0.04 ^c
Potassium	0.71 ± 0.00 ^a	0.55 ± 0.02 ^b	0.43 ± 0.01 ^c
Phosphorous	7.03 ± 0.03 ^a	2.74 ± 0.02 ^b	6.45 ± 0.07 ^c
<i>Microminerals (µg/g)</i>			
Iron	28.30 ± 0.11 ^a	28.73 ± 0.08 ^b	40.36 ± 0.55 ^c
Copper	8.93 ± 0.03 ^a	7.46 ± 0.09 ^b	6.21 ± 0.32 ^c
Manganese	1.64 ± 0.02 ^a	2.64 ± 0.02 ^b	2.84 ± 0.02 ^c
Zinc	24.03 ± 0.45 ^a	24.61 ± 0.19 ^b	54.46 ± 0.17 ^c

Values are mean% ± SE. Values in the same row followed by different letters are significantly different ($p < 0.05$).

TABLE 2: Fatty acid composition of three species of Bagridae fish.

Fatty acid	<i>Hemibagrusnemurus</i>	<i>Hemibagruswyckii</i>	<i>Mystusnigriceps</i>
C12:0 (lauric acid)	1.97 ± 0.05 ^a	3.94 ± 0.01 ^b	4.27 ± 0.01 ^c
C14:0 (myristic acid)	2.38 ± 0.03 ^a	2.72 ± 0.08 ^b	2.89 ± 0.01 ^c
C16:0 (palmitic acid)	27.23 ± 0.06 ^a	30.70 ± 0.06 ^b	25.59 ± 0.01 ^c
C18:0 (stearic acid)	16.17 ± 0.02 ^a	13.54 ± 0.02 ^b	10.64 ± 0.01 ^c
C20:0 (arachidic acid)	0.23 ± 0.01 ^a	0.14 ± 0.01 ^b	0.37 ± 0.01 ^c
C16:1 (palmitoleic acid)	1.78 ± 0.00 ^a	1.27 ± 0.04 ^b	4.93 ± 0.01 ^c
C18:1 (oleic acid)	16.86 ± 0.01 ^a	16.18 ± 0.01 ^b	16.97 ± 0.01 ^c
C18:2 (linoleic acid)	3.24 ± 0.02 ^a	3.84 ± 0.33 ^b	4.55 ± 0.001 ^c
C18:3 (linolenic acid)	1.12 ± 0.01 ^a	0.82 ± 0.01 ^b	1.57 ± 0.00 ^c
C20:4 (arachidonic acid)	12.13 ± 0.01 ^a	12.49 ± 0.09 ^b	12.41 ± 0.02 ^c
C20:5 (eicosapentaenoic acid; EPA)	2.06 ± 0.08 ^a	2.34 ± 0.04 ^b	1.35 ± 0.01 ^c
C22:6 (docosahexaenoic acid; DHA)	15.65 ± 0.01 ^a	11.56 ± 0.04 ^b	13.64 ± 0.02 ^c

Values are mean% ± SE. Values in the same row with different superscripts are significantly different ($p < 0.05$).

1:0.97. It was found that all three fish species had ω -6/ ω -3 ratios within the recommended range for a typical Indonesian diet. In contrast, a typical Western diet is characterized by a high ω -6/ ω -3 ratio of 10:1 to 20:1, while a Mediterranean diet has a ratio of 4:1 to 5:1 [43]. Additionally, the P/S ratio analysis showed that the three Bagridae fish species are sources of PUFAs that meet the requirements for food safety.

The Food and Agriculture Organization (FAO) and the World Health Organization (WHO) have recommended AI and TI values ranging from 0.4 to 0.5 for promoting human health [44]. In the current study, the observed AI values in the three freshwater species of Bagridae ranged from 0.73 to 0.99, while the TI values varied between 0.40 and 0.63. This finding is associated with a notable disparity

TABLE 3: Fatty acid ω -6/ ω -3 and polyunsaturated/saturated fatty acid ratio of three species of Bagridae fish.

	<i>Hemibagrusnemurus</i>	<i>Hemibagruswyckii</i>	<i>Mystusnigriceps</i>
ω -6/ ω -3 ratio	0.82 ± 0.00 ^a	1.11 ± 0.02 ^b	1.02 ± 0.01 ^c
P/S ratio	0.69 ± 0.01 ^a	0.61 ± 0.06 ^b	0.76 ± 0.00 ^c
AI	0.87 ± 0.01 ^a	0.99 ± 0.06 ^b	0.73 ± 0.00 ^c
TI	0.40 ± 0.01 ^a	0.63 ± 0.01 ^b	0.43 ± 0.00 ^c

P/S: polyunsaturated/saturated fatty acid; AI: atherogenic index; TI: thrombogenic index. Values are mean% ± SE. Values in the same row followed by different letters are significantly different ($p < 0.05$).

TABLE 4: Amino acid composition of three species of Bagridae fish.

Amino acid	<i>Hemibagrusnemurus</i>	<i>Hemibagruswyckii</i>	<i>Mystusnigriceps</i>
Aspartic acid	10.39 ± 0.01 ^a	9.61 ± 0.04 ^b	11.28 ± 0.02 ^c
Glutamic acid	24.34 ± 0.00 ^a	21.58 ± 0.41 ^b	19.29 ± 0.03 ^c
Serine	3.74 ± 0.00 ^a	5.40 ± 0.03 ^b	7.50 ± 0.06 ^c
Glycine	4.02 ± 0.00 ^a	5.77 ± 0.04 ^b	4.69 ± 0.02 ^c
Histidine	2.39 ± 0.00 ^a	2.90 ± 0.03 ^b	2.21 ± 0.02 ^c
Arginine	6.17 ± 0.01 ^a	8.27 ± 0.03 ^b	6.22 ± 0.03 ^c
Threonine	4.49 ± 0.01 ^a	4.19 ± 0.03 ^b	4.39 ± 0.04 ^c
Alanine	7.03 ± 0.00 ^a	6.04 ± 0.02 ^b	7.89 ± 0.01 ^c
Proline	2.40 ± 0.00 ^a	2.71 ± 0.01 ^b	2.21 ± 0.02 ^c
Tyrosine	3.91 ± 0.01 ^a	3.29 ± 0.06 ^b	3.82 ± 0.04 ^c
Valine	3.73 ± 0.00 ^a	4.02 ± 0.01 ^b	3.85 ± 0.04 ^c
Methionine	3.42 ± 0.00 ^a	1.96 ± 0.02 ^b	1.27 ± 0.03 ^c
Cystine	0.87 ± 0.00 ^a	0.76 ± 0.02 ^b	0.75 ± 0.03 ^c
Isoleucine	3.82 ± 0.001 ^a	3.96 ± 0.003 ^b	3.66 ± 0.002 ^c
Leucine	6.15 ± 0.000 ^a	6.69 ± 0.002 ^b	6.28 ± 0.004 ^c
Phenylalanine	5.13 ± 0.000 ^a	6.03 ± 0.005 ^b	5.18 ± 0.027 ^c
Lysine	9.61 ± 0.001 ^a	9.82 ± 0.026 ^b	9.87 ± 0.003 ^c

Values are mean% ± SE. Values in the same row followed by different letters are significantly different ($p < 0.05$).

in the values of SFAs among the three Bagridae fish species. Based on these findings, we confirm that consuming the meat of the three freshwater species of Bagridae fish caught in the wild does not endanger consumer health and can serve as a suitable source of food safety and nutritional quality.

Fish with a high crude protein content, as shown by the three studied Bagridae fish (Table 1), are regarded as being nutritionally useful for human health. The protein and amino acid contents can directly indicate the nutritional quality of meat [45]. As shown in Table 4, *H. wyckii* had a higher content of essential amino acids (EAAs) than *H. nemurus* and *M. nigriceps*. Among the EAAs, lysine (Lys) was found to be present in the highest amount in all three species. Lys is necessary for tissue growth and repair, reduced stress-induced anxiety, and the conversion of fatty acids into energy [46–48].

More than 60% of the amino acid composition in all three Brigade fish studied comes from nonessential amino acids (NEAAs), particularly glutamic acid (Table 4). Glutamic acid and its amine, glutamine, are highly abundant amino acids found in fish in the free and protein-bound

forms [49]. Other amino acids found at abundant levels in the three fish also play important physiological and biochemical functions. The NEAA arginine, for example, is involved in protein synthesis, creatine formation, and various other biochemical processes within the body. The EAAs leucine, phenylalanine, and methionine play important roles in the synthesis of nitric oxide, a molecule that can expand blood vessels [49].

5. Conclusions

The present study provides insightful information about the fatty acid and amino acid compositions, biometric measurements, proximate analysis, and mineral composition of three local freshwater species Bagridae fish in the Kampar Kanan river, Indonesian. The findings show their potential as a nutrient-dense, sustainable food source that can support efforts to increase food security. These fish species might be incorporated into the local diet to increase nutrient intake and alleviate nutritional inadequacies, which would ultimately improve Indonesia's overall food security status.

Data Availability

The data utilized in this research has not been previously released or published in any form. The datasets employed and/or analysed during the present study can be obtained by contacting the corresponding author.

Conflicts of Interest

The authors confirm that they do not have any conflicting interests.

Acknowledgments

The authors acknowledge and express their gratitude for the financial support received from the Ministry of Education, Culture, Research, and Technology of the Republic of Indonesia, specifically through grant no. 076/E5/PG. 02.00/PT/2022, which funded the research. Furthermore, the authors extend their appreciation to the students and fishers involved in the research conducted in the upper reaches of the Kampar Kanan River, Indonesia.

References

- [1] A. E. Hasselberg, I. Aakre, J. Scholtens et al., "Fish for food and nutrition security in Ghana: challenges and opportunities," *Global Food Security*, vol. 26, article 100380, 2020.
- [2] N. Tran, U. P. Rodriguez, Y. Chan, Y. M. Auang, L. Chu, and A. H. M. S. Islam, "Future scenarios of fish supply and demand for food and nutrition security in Bangladesh: an analysis with the AsiaFish model," *Aquaculture*, vol. 568, article 739288, 2023.
- [3] C. Z. Levkoe, K. Lowitt, and C. Nelson, "Fish as food: exploring a food sovereignty approach to small-scale fisheries," *Marine Policy*, vol. 85, pp. 65–70, 2017.
- [4] M. Ravi, N. S. Hettiarachchy, V. Ganesh, and S. Ricke, "Enhancement of antimicrobial activities of naturally occurring phenolic compounds by nanoscale delivery against *Listeria monocytogenes*, *Escherichia coli* O157: H7 and *Salmonella typhimurium* in broth and chicken meat system," *Journal of Food Safety*, vol. 31, no. 4, pp. 462–471, 2011.
- [5] N. Ariani, I. Suharman, H. Syandri, and A. Mardiah, "Diversity and distribution of fish fauna of upstream and downstream areas at Koto Panjang Reservoir, Riau Province, Indonesia," *F1000 Research*, vol. 8, no. 1435, pp. 1–16, 2019.
- [6] D. Syafrialdi, D. I. Roesma, and H. Syandri, "Length-weight relationships and condition factor of two-spot catfish (*Mystus nigriceps* [Valenciennes, 1840]) (Pisces, Bagridae), from Kampar Kanan River and Kampar Kiri River in Indonesia," *Pakistan Journal of Biological Science*, vol. 23, no. 12, pp. 1636–1642, 2020.
- [7] N. Ariani, "Native species in Kampar Kanan River, Riau Province Indonesia," *International Journal of Fisheries and Aquatic Studies*, vol. 2, no. 5, pp. 213–217, 2015.
- [8] A. Azrita, H. Syandri, N. Ariani, A. Mardiah, and I. Suharman, "The utilization of new products formulated from water coconut, palm sap sugar, and fungus to increase nutritional feed quality, feed efficiency, growth, and carcass of gurami sago (*Osphronemus goramy* Lacepede, 1801) juvenile," *F1000 Research*, vol. 10, no. 1121, pp. 1–16, 2021.
- [9] C. Bene, R. Arthur, H. Norbury et al., "Contribution of fisheries and aquaculture to food security and poverty reduction: assessing the current evidence," *World Development*, vol. 79, pp. 177–196, 2016.
- [10] N. Ariyani, I. Suharman, S. Hasibuan, N. Asia, and H. Syandri, "Fatty acid composition on diet and carcasses, growth, body indices and profile serum of Asian redtail catfish (*Hemibagrus nemurus*) fed a diet containing different levels of EPA and DHA," *F1000 Research*, vol. 11, no. 1409, pp. 1–23, 2022.
- [11] E. N. Kingsley, O. O. Cyril, and O. I. Patience, "Potential contribution of selected wild fish species to the minerals intake of pregnant and lactating women, children and adults in rural riverine communities of Edo State: insights and outcomes," *Measurement: Food*, vol. 8, article 100063, pp. 1–7, 2022.
- [12] A. A. Sausa, L. Nora, D. L. A. Lopes et al., "Vegetable choline in feed for Nile tilapia (*Oreochromis niloticus*) raised in a biofloc technology system (BFT): biofloc composition, chemical composition, and fatty acid profiles in meat," *Aquaculture*, vol. 545, article 737174, 2021.
- [13] A. M. Eltweri, A. L. Thomas, M. Metcalfe, P. C. Calder, A. R. Dennis, and D. J. Bowrey, "Potential applications of fish oils rich in omega-3 polyunsaturated fatty acids in the management of gastrointestinal cancer," *Clinical Nutrition*, vol. 36, no. 1, pp. 65–78, 2017.
- [14] N. Tintle, T. Rice, I. Cheng et al., "The association between genetically elevated polyunsaturated fatty acids and risk of cancer," *eBioMedicine*, vol. 91, article 10210, 2023.
- [15] E. K. Aglago, I. Huybrechts, N. Murphy et al., "Consumption of fish and long-chain n-3 polyunsaturated fatty acids is associated with reduced risk of colorectal cancer in a large European cohort," *Clinical Gastroenterology and Hepatology*, vol. 18, no. 3, pp. 654–666.e6, 2020.
- [16] H. Jiang, X. Shi, T. Y. Fan et al., "Dietary omega-3 polyunsaturated fatty acids and fish intake and risk of age-related macular degeneration," *Clinical Nutrition*, vol. 40, no. 12, pp. 5662–5673, 2021.
- [17] M. A. Rincón-Cervera, N. V. González-Barriga, R. Valenzuela, S. Lopez-Arana, J. Romero, and A. Valenzuela, "Profile and distribution of fatty acids in edible parts of commonly consumed marine fishes in Chile," *Food Chemistry*, vol. 274, pp. 123–129, 2019.
- [18] S. Sun, T. Ren, X. Li, X. Cao, and J. Gao, "Polyunsaturated fatty acids synthesized by freshwater fish: a new insight to the roles of *elov12* and *elov15* in vivo," *Biochemical and Biophysical Research Communications*, vol. 532, no. 3, pp. 414–419, 2020.
- [19] M. B. Behyar, M. Hasanzadeh, F. Seidi, and N. Sadjou, "Sensing of amino acids: critical role of nanomaterials for the efficient biomedical analysis," *Microchemical Journal*, vol. 188, article 108452, 2023.
- [20] P. Shi, J. Xu, Y. Wang, S. Xu, and X. Yan, "Eicosapentaenoic acid mitigates palmitic acid-induced heat shock response, inflammation and repair processes in fish intestine," *Fish & Shellfish Immunology*, vol. 124, pp. 362–371, 2022.
- [21] J. Wen, L. Zeng, Y. Xu, Y. Sun, Z. Chen, and S. Fan, "Proximate composition, amino acid and fatty acid composition of fish maws," *Natural Product Research*, vol. 30, no. 2, pp. 214–217, 2016.
- [22] F. O. S. Duarte, F. G. de Paula, C. S. Prado et al., "Better fatty acids profile in filets of Nile tilapia (*Oreochromis niloticus*) supplemented with fish oil," *Aquaculture*, vol. 534, article 736241, 2021.

- [23] K. L. Weaver, P. Ivester, J. A. Chilton, M. D. Wilson, P. Pandey, and F. H. Chilton, "The content of favorable and unfavorable polyunsaturated fatty acids found in commonly eaten fish," *Journal of the American Dietetic Association*, vol. 108, no. 7, pp. 1178–1185, 2008.
- [24] AOAC, *Official Methods of Analysis*, Association of Official Analytical Chemists (AOAC), Washington, DC., USA, 15th edition, 1990.
- [25] S. A. Cohen, "Amino acid analysis using pre-column derivatization with 6-aminoquinolyl-N-hydroxysuccinimidyl carbamate," *Amino Acid Analysis Protocols*, vol. 211, pp. 143–154, 2000.
- [26] M. A. Rajion, J. G. McLean, and R. N. Cahill, "Essential fatty acid metabolism in the fetal and neonatal lamb," *Australian Journal of Biological Sciences*, vol. 38, no. 1, pp. 33–40, 1985.
- [27] T. L. Ulbricht and D. A. Southgate, "Coronary heart disease: seven dietary factors," *The Lancet*, vol. 338, no. 8773, pp. 985–992, 1991.
- [28] D. B. Duncan, "Multiple range and multiple F tests," *Biometrics*, vol. 11, no. 1, pp. 1–42, 1955.
- [29] R. Froese, "Cube law, condition factor and weight-length relationships: history, meta-analysis and recommendations," *Journal of Applied Ichthyology*, vol. 22, no. 4, pp. 241–253, 2006.
- [30] O. O. Famoofo and W. O. Abdul, "Biometry, condition factors and length-weight relationships of sixteen fish species in Iwopin fresh-water ecotype of Lekki Lagoon, Ogun State, southwest Nigeria," *Heliyon*, vol. 6, no. 1, article e02957, 2020.
- [31] I. B. Bagenal and F. W. Tesch, "Age and growth," in *Methods for Assessment of Fish Production in Fresh Waters*, T. Bagenal, Ed., pp. 101–136, Blackwell Science Publications, 1978.
- [32] E. O. Ahmed, M. E. Ali, and A. A. Aziz, "Length-weight relationships and condition factors of six fish species in Atbara River and Khashm El-Girba reservoir, Sudan," *International Journal of Agriculture Sciences*, vol. 3, no. 1, pp. 65–70, 2011.
- [33] E. Ujjania, M. P. S. Kohli, and L. L. Sharma, "Length-weight relationship and condition factors of Indian major carps (*C. catla*, *L. rohita* and *C. mrigala*) in Mahi Bajaj Sagar, India," *Research Journal of Biology*, vol. 2, no. 1, pp. 30–36, 2012.
- [34] J. P. Bonjour, "Calcium and phosphate: a duet of ions playing for bone health," *Journal of the American College of Nutrition*, vol. 30, Supplement 5, pp. 438S–448S, 2011.
- [35] B. P. Mohanty, T. V. Sankar, S. Ganguly et al., "Micronutrient composition of 35 food fishes from India and their significance in human nutrition," *Biological Trace Element Research*, vol. 174, no. 2, pp. 448–458, 2016.
- [36] H. Syandri, A. Azrita, A. Mardi, N. Aryani, and A. Dihami, "The proximate composition, amino acid profile, fatty acid content, and mineral content of scale flour from three fish species as potential feeds for fish fry," *F1000 Research*, vol. 12, no. 1144, pp. 1–19, 2023.
- [37] F. Jabeen and A. A. Chudhry, "Chemical compositions and fatty acid profiles of three freshwater fish species," *Food Chemistry*, vol. 125, no. 3, pp. 991–996, 2011.
- [38] A. A.URNADIA, A. Azrina, and I. Amin, "Proximate composition and energetic value of selected marine fish and shellfish from the west coast of peninsular Malaysia," *International Food Research Journal*, vol. 18, pp. 137–148, 2011.
- [39] S. C. Dyal, L. Balas, N. G. Bazan et al., "Polyunsaturated fatty acids and fatty acid-derived lipid mediators: recent advances in the understanding of their biosynthesis, structures, and functions," *Progress in Lipid Research*, vol. 86, article 101165, 2022.
- [40] N. Kim, U. D. Sohn, V. Mangannan et al., "Leukotrienes in acetylcholine-induced contraction of esophageal circular smooth muscle in experimental esophagitis," *Gastroenterology*, vol. 112, no. 5, pp. 1548–1558, 1997.
- [41] D. Swanson, R. Block, and S. A. Mousa, "Omega-3 fatty acids EPA and DHA: health benefits throughout life," *Advances in Nutrition*, vol. 3, no. 1, pp. 1–7, 2012.
- [42] R. P. Mason, S. C. R. Sherratt, and L. H. Eckel, "Omega-3-fatty acids: do they prevent cardiovascular disease?," *Best Practice & Research Clinical Endocrinology & Metabolism*, vol. 37, no. 3, article 101681, 2023.
- [43] J. M. Seddon, J. Cote, N. Davis, and B. Rosner, "Progression of age-related macular degeneration," *Archives of Ophthalmology*, vol. 121, no. 4, pp. 785–792, 2003.
- [44] FAO/WHO, *Fat and Oils in Human Nutrition: Report of a Joint Expert Consultation*, FAO/WHO, Rome, Italy, 1994.
- [45] S. M. Pawar and S. R. Sonaware, "Fish muscle protein highest source of energy," *International Journal of Biodiversity and Conservation*, vol. 5, no. 7, pp. 433–435, 2013.
- [46] D. Tomé and C. Bos, "Lysine requirement through the human life cycle," *The Journal of Nutrition*, vol. 137, no. 6, pp. 1642S–1645S, 2007.
- [47] M. Watford, "Glutamine and glutamate: nonessential or essential amino acids?," *Animal Nutrition*, vol. 1, no. 3, pp. 119–122, 2015.
- [48] X. Li, S. Zheng, and G. Wu, "Nutrition and metabolism of glutamate and glutamine in fish," *Amino Acids*, vol. 52, no. 5, pp. 671–691, 2020.
- [49] M. Li, Y. Wu, and L. Ye, "The role of amino acids in endothelial biology and function," *Cells*, vol. 11, no. 8, p. 1372, 2022.

Analysis of Fatty Acids and Amino Acids of Three Local Freshwater Bagridae Fish Species In The Kampar Kanan River Indonesia For Food Security

ORIGINALITY REPORT

16%

SIMILARITY INDEX

16%

INTERNET SOURCES

%

PUBLICATIONS

%

STUDENT PAPERS

PRIMARY SOURCES

1	jfmr.ub.ac.id Internet Source	<1 %
2	libinfo.uark.edu Internet Source	<1 %
3	repositorio.unesp.br Internet Source	<1 %
4	www.iarc.who.int Internet Source	<1 %
5	accounts.public.ce.basespace.illumina.com Internet Source	<1 %
6	medwinpublishers.com Internet Source	<1 %
7	pure.roehampton.ac.uk Internet Source	<1 %
8	repositorio.ufc.br Internet Source	<1 %

9	Internet Source	<1 %
10	journal.ipb.ac.id Internet Source	<1 %
11	summit.sfu.ca Internet Source	<1 %
12	www.oecd-ilibrary.org Internet Source	<1 %
13	cgspace.cgiar.org Internet Source	<1 %
14	eprints.ncl.ac.uk Internet Source	<1 %
15	moam.info Internet Source	<1 %
16	eprints.kaums.ac.ir Internet Source	<1 %
17	par.nsf.gov Internet Source	<1 %
18	jtrolis.ub.ac.id Internet Source	<1 %
19	ppgca.evz.ufg.br Internet Source	<1 %
20	vivo.brown.edu Internet Source	<1 %

21	www.academicjournals.org Internet Source	<1 %
22	www.rjd.miami.edu Internet Source	<1 %
23	civilejournal.org Internet Source	<1 %
24	neurotree.org Internet Source	<1 %
25	pure.johnshopkins.edu Internet Source	<1 %
26	www.sun.ac.za Internet Source	<1 %
27	dspace.uef.fi Internet Source	<1 %
28	www.thefreelibrary.com Internet Source	<1 %
29	dx.doi.org Internet Source	<1 %
30	www.researchwithnj.com Internet Source	<1 %
31	www.slideshare.net Internet Source	<1 %
32	ciari.icar.gov.in Internet Source	<1 %

33	journalspress.com Internet Source	<1 %
34	terkko.helsinki.fi Internet Source	<1 %
35	www.intechopen.com Internet Source	<1 %
36	www.sciepub.com Internet Source	<1 %
37	www.thermofisher.com Internet Source	<1 %
38	findresearcher.sdu.dk Internet Source	<1 %
39	seer.sct.embrapa.br Internet Source	<1 %
40	www.hrpub.org Internet Source	<1 %
41	www.naturalproductsinsider.com Internet Source	<1 %
42	www.rural21.com Internet Source	<1 %
43	www.sci-int.com Internet Source	<1 %
44	journals.lww.com Internet Source	<1 %

45	www.j3.jstage.jst.go.jp Internet Source	<1 %
46	www.koreascience.kr Internet Source	<1 %
47	www.jbc.org Internet Source	<1 %
48	www.tara.tcd.ie Internet Source	<1 %
49	static.sites.s bq.org.br Internet Source	<1 %
50	www.coursehero.com Internet Source	<1 %
51	www.sciencedirect.com Internet Source	<1 %
52	health-and-age.squarespace.com Internet Source	<1 %
53	knapublishing.com Internet Source	<1 %
54	acs.agr.hr Internet Source	<1 %
55	vdocuments.com.br Internet Source	<1 %
56	web.unbc.ca Internet Source	<1 %

57 worldwidescience.org <1 %
Internet Source

58 eprints.ugd.edu.mk <1 %
Internet Source

59 ojs.unud.ac.id <1 %
Internet Source

60 orca.cardiff.ac.uk <1 %
Internet Source

61 www.garj.org <1 %
Internet Source

62 zoe.com <1 %
Internet Source

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off